

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
14 July 2005 (14.07.2005)

PCT

(10) International Publication Number  
**WO 2005/063254 A2**

(51) International Patent Classification<sup>7</sup>: **A61K 31/551**,  
A61P 25/28, C07D 403/04

(21) International Application Number:  
PCT/US2004/043224

(22) International Filing Date:  
21 December 2004 (21.12.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/531,927 22 December 2003 (22.12.2003) US  
60/548,090 24 February 2004 (24.02.2004) US  
60/548,604 27 February 2004 (27.02.2004) US

(71) Applicant (for all designated States except US): **ACADIA PHARMACEUTICALS INC.** [US/US]; 3911 Sorrento Valley Road, San Diego, California 92121 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **EK, Fredrik** [SE/SE]; Dag Hammarskjöldsv. 55, S-224 64 Lund (SE). **OLSSON, Roger** [SE/SE]; Klagshamnsgatan 99A, S-23044 Bunkeflo Strand (SE). **OHLSSON, Jörgen** [SE/SE]; Jägaregatan 44, S-226 53 Lund (SE).

(74) Agent: **HART, Daniel**; KNOBBE MARTENS OLSON & BEAR LLP, 2040 Main Street, 14th Floor, Irvine, California 92614 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**

— of inventorship (Rule 4.17(iv)) for US only

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: AMINO SUBSTITUTED DIARYL[A,D]CYCLOHEPTENE ANALOGS AS MUSCARINIC AGONISTS AND METHODS OF TREATMENT OF NEUROPSYCHIATRIC DISORDERS

(57) Abstract: Disclosed herein are analogs of clozapine and pharmaceutically acceptable salts, esters, amides, or prodrugs thereof; methods of synthesizing the analogs; and methods of using the analogs for treating neuropsychiatric disorders. In some embodiments, the analogs are amino substituted diaryl[a,d]cycloheptenes.



WO 2005/063254 A2

**AMINO SUBSTITUTED DIARYL[a,d]CYCLOHEPTENE ANALOGS AS**  
**MUSCARINIC AGONISTS AND METHODS OF TREATMENT OF**  
**NEUROPSYCHIATRIC DISORDERS**

Background of the Invention

Field of the Invention

[0001] Certain aspects of the present disclosure relate to methods for treatment of neuropsychiatric disorders, pain and other disorders by compounds that modulate the activity of muscarinic receptors, in particular the subtypes M1, thereby modulating neuronal activities associated with the development of neuropsychiatric disorders. Aspects of the invention also relate to compounds that selectively interact with this receptor subtype and methods of identifying said compounds.

Description of the Related Art

[0002] Muscarinic cholinergic receptors mediate the actions of the neurotransmitter acetylcholine in the central and peripheral nervous systems, gastrointestinal system, heart, endocrine glands, lungs, and other tissues. Muscarinic receptors play a central role in the central nervous system for higher cognitive functions, as well as in the peripheral parasympathetic nervous system. Five distinct muscarinic receptor subtypes have been identified, m1-m5. The m1 subtype is the predominant subtype found in the cerebral cortex and is believed to be involved in the control of cognitive functions; m2 is the predominant subtype found in heart and is believed to be involved in the control of heart rate; m3 is believed to be involved in gastrointestinal and urinary tract stimulation as well as sweating and salivation; m4 is present in brain and may be involved in locomotion; and m5, present in brain, may be involved in certain functions of the central nervous system associated with the dopaminergic system.

[0003] Conditions associated with cognitive impairment, such as Alzheimer's disease, are accompanied by loss of acetylcholine in the brain. This is believed to be the result of degeneration of cholinergic neurons in the basal forebrain, which innervate areas of the association cortex, and hippocampus, which is involved in higher processes.

[0004] Efforts to increase acetylcholine levels have focused on increasing levels of choline, the precursor for acetylcholine synthesis, and on blocking acetylcholine esterase (AChE), the enzyme that metabolizes acetylcholine. Administration of choline or phosphatidylcholine has not been very successful. AChE inhibitors have shown some




therapeutic efficacy, but may cause cholinergic side effects due to peripheral acetylcholine stimulation, including abdominal cramps, nausea, vomiting, diarrhea, anorexia, weight loss, myopathy and depression. Gastrointestinal side effects have been observed in about a third of the patients treated. In addition, some AChE inhibitors, such as tacrine, have also been found to cause significant hepatotoxicity, with elevated liver transaminases observed in about 30% of patients. The adverse effects of AChE inhibitors have limited their clinical utility.

[0005] Known m1 muscarinic agonists such as arecoline have also been found to be weak agonists of m2 as well as m3 subtype and are not very effective in treating cognitive impairment, most likely because of dose-limiting side effects.


[0006] There is a need for compounds that increase acetylcholine signaling or effect in the brain. Specifically there is a need for muscarinic agonists that are active at various muscarinic receptor subtypes in the central and peripheral nervous system. Furthermore, there is a need for more highly selective muscarinic agonists, such as m1- or m4-selective agents, both as pharmacological tools and as therapeutic agents.

#### Summary of the Invention

[0007] Disclosed herein is a compound of Formula I, II, or XV:




or a pharmaceutically acceptable salt, ester, amide, or prodrug thereof, wherein A is selected from the group consisting of




X is nitrogen, CH, or CH<sub>2</sub>; X' is C or CH, wherein when X' is C, there is a double bond between X and X' and wherein when X' is CH, there is a single bond between X and X'; each Y is separately selected from the group consisting of nitrogen, oxygen, or CH; each W is separately selected from the group consisting of nitrogen, CH, oxygen, or sulfur; each n is separately selected from the group consisting of 0, 1, 2, 3, and 4; m is selected from the group consisting of 1, 2, and 3; each R<sub>1</sub> is separately absent or is separately selected from the group consisting of hydrogen, halogen, amine, optionally substituted C<sub>1-20</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-20</sub> alkenyl, optionally substituted C<sub>2-20</sub> alkynyl, optionally substituted C<sub>1-20</sub> -alkoxyalkyl, and optionally substituted aryl and arylalkyl; L is absent or is selected from the group consisting of -NH(CH<sub>2</sub>)<sub>n</sub>- and -(CH<sub>2</sub>)<sub>n</sub>-; a, b, c, and d are each independently selected from the group consisting of carbon, nitrogen, oxygen, and sulfur, or each is independently absent, provided that at least three of a, b, c, or d are present, provided that at least one of a, b, c, or d is carbon, and provided that no two adjacent a, b, c, or d are both oxygen or both sulfur; e, f, g, and h are each independently selected from the group consisting of carbon, nitrogen, oxygen, and sulfur, or each is independently absent, provided that at least three of e, f, g, or h are present, provided that at least one of e, f, g, or h is carbon, and provided that no two adjacent e, f, g, or h are both oxygen or both sulfur; R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, and R<sub>5</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>, or R<sub>2</sub> and R<sub>3</sub>, or R<sub>3</sub> and R<sub>4</sub>, or R<sub>4</sub> and R<sub>5</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety; R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, and R<sub>9</sub>, are each independently selected from the

group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>, or R<sub>6</sub> and R<sub>7</sub>, or R<sub>7</sub> and R<sub>8</sub>, or R<sub>8</sub> and R<sub>9</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety; Z is selected from the group consisting of NR<sub>11</sub>, oxygen, sulfur, and CH<sub>2</sub>; R<sub>10</sub> is selected from the group consisting of hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl optionally substituted aryl, optionally substituted arylalkyl, and perhaloalkyl; and R<sub>11</sub> is selected from the group consisting of hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, and optionally substituted arylalkyl; R<sub>12</sub> and R<sub>13</sub> are separately selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>, or R<sub>12</sub> and R<sub>13</sub>, taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety; and any bond represented by a dashed and solid line represents a bond selected from the group consisting of a carbon-carbon single bond and a carbon-carbon double bond; provided that the compound of Formulae I or XV are not clozapine or N-desmethylozapine.

[0008] In some embodiments, the compound has a structure set forth in Formulas III or IV.





(III)









(IV)

[0009] In some embodiments, the compound is selected from the group consisting of:



[0010] In some embodiments, the compound is selected from the group consisting of:




**[0010]** In some embodiments, none of a, b, c, or d is absent. In some embodiments, none of e, f, g, or h is absent. In some embodiments a, b, c, and d are carbon. In some embodiments e, f, g, and h are carbon. In some embodiments, R<sub>2</sub> is selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, and optionally substituted C<sub>1-6</sub> alkyloxy. In some embodiments, the alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. In some embodiments, the alkyloxy is selected from the group consisting of methoxy, ethoxy, propoxy, isopropoxy, butoxy, sec-butoxy, and tert-butoxy. In some embodiments, the halogen is selected from the group consisting of fluoro, chloro, and bromo. In some embodiments, R<sub>2</sub> is selected from the group consisting of hydrogen, methyl, methoxy, and chloro. In some embodiments, R<sub>3</sub> is selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, and NO<sub>2</sub>. In some embodiments, the alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. In some embodiments, the alkyloxy is selected from the group consisting of methoxy, ethoxy, propoxy, isopropoxy, butoxy, sec-butoxy, and tert-butoxy. In some embodiments, the halogen is selected from the group consisting of chloro, bromo, and iodo. In some embodiments, R<sub>3</sub> is selected from the group consisting of hydrogen, methyl, methoxy, chloro, bromo, iodo, and NO<sub>2</sub>. In some embodiments, R<sub>4</sub> is selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, perhaloalkyl, SO<sub>2</sub>R<sub>10</sub>, and NO<sub>2</sub>. In some embodiments, the alkyl is selected from the group

consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. In some embodiments, the perhaloalkyl is perfluoroalkyl. In some embodiments, the perfluoroalkyl is trifluoromethyl. In some embodiments, the halogen is selected from the group consisting of fluoro, chloro, and bromo. In some embodiments,  $R_{10}$  is hydrogen or optionally substituted  $C_{1-6}$  alkyl, wherein in some embodiments, the alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. In some embodiments,  $R_4$  is selected from the group consisting of hydrogen, methyl, fluoro, chloro, bromo, trifluoromethyl,  $SO_2CH_3$ , and  $NO_2$ . In some embodiments,  $R_5$  is selected from the group consisting of hydrogen, halogen, and optionally substituted  $C_{1-6}$  alkyl, wherein in some embodiments, the alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl and wherein in some embodiments, the halogen is selected from the group consisting of fluoro, chloro, and bromo. In some embodiments,  $R_5$  is hydrogen or chloro. In some embodiments,  $R_6$  is hydrogen or optionally substituted  $C_{1-6}$  alkyl. In some embodiments,  $R_6$  is hydrogen. In some embodiments,  $R_7$  is selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, perhaloalkyl, CN,  $SO_2R_{10}$ , and  $NO_2$ , wherein in some embodiments, the alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl; wherein in some embodiments, the halogen is selected from the group consisting of fluoro, chloro, and bromo; wherein in some embodiments, perhaloalkyl is perfluoroalkyl; wherein in some embodiments, perfluoroalkyl is trifluoromethyl. In some embodiments,  $R_{10}$  is hydrogen or optionally substituted  $C_{1-6}$  alkyl, wherein in some embodiments, the alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. In some embodiments,  $R_7$  is selected from the group consisting of hydrogen, methyl, chloro, trifluoromethyl,  $SO_2CH_3$ , CN, and  $NO_2$ . In some embodiments,  $R_8$  is selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, wherein in some embodiments, the alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl and wherein in some embodiments, the halogen is selected from the group consisting of fluoro, chloro, and bromo. In some embodiments,  $R_8$  is selected from the group consisting of hydrogen, chloro, and bromo. In some embodiments,  $R_9$  is selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, and perhaloalkyl; wherein in some embodiments, the alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl; wherein in some embodiments, the halogen is selected from the group consisting





of fluoro, chloro, and bromo; wherein in some embodiments, perhaloalkyl is perfluoroalkyl; wherein in some embodiments, perfluoroalkyl is trifluoromethyl. In some embodiments,  $R_9$  is selected from the group consisting of hydrogen, chloro, methyl, and trifluoromethyl. In some embodiments,  $R_1$  is selected from the group consisting of hydrogen, optionally substituted  $C_{1-6}$  alkyl, and optionally substituted aryl, wherein in some embodiments, the alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. In some embodiments,  $R_1$  is hydrogen. In some embodiments,  $X$  is nitrogen. In some embodiments,  $Y$  is  $NH$ . In some embodiments,  $L$  is absent or is selected from the group consisting of  $-NHCH_2-$ ,  $-NH-$ , and  $-CH_2-$ . In some embodiments,  $A$  is selected from the group consisting of:




and  $n$  is selected from the group consisting of 0, 1, and 2.


[0011] Also disclosed herein is a method of synthesizing a compound of Formula V or VI,




comprising reacting a compound of Formula VII




with a compound of Formula VIII



to form a fused ring compound of Formula IX,




and reacting the compound of Formula IX with a compound of Formula X




to obtain a compound of Formula V, wherein X is a halogen; R<sub>1</sub> is selected from the group consisting of hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub> -alkoxyalkyl, and optionally substituted aryl and arylalkyl; R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, and R<sub>5</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>, or R<sub>2</sub> and R<sub>3</sub>, or R<sub>3</sub> and R<sub>4</sub>, or R<sub>4</sub> and R<sub>5</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety; R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, and R<sub>9</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>, or R<sub>6</sub> and R<sub>7</sub>, or R<sub>7</sub> and R<sub>8</sub>, or R<sub>8</sub> and R<sub>9</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety.


[0012] Also disclosed herein is a combinatorial library of at least 220 dibenzo[*b,e*][1,4]diazepine[*a,d*]cycloheptene compounds that can be formed by reacting a compound of Formula VII,



with a compound of Formula VIII and




a compound of Formula X,




wherein X is a halogen; R<sub>1</sub> is selected from the group consisting of hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub> -alkoxyalkyl, and optionally substituted aryl and arylalkyl; R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, and R<sub>5</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>, or R<sub>2</sub> and R<sub>3</sub>, or R<sub>3</sub> and R<sub>4</sub>, or R<sub>4</sub> and R<sub>5</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety; R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, and R<sub>9</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>, or R<sub>6</sub> and R<sub>7</sub>, or R<sub>7</sub> and R<sub>8</sub>, or R<sub>8</sub> and R<sub>9</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety.


[0013] Also disclosed herein is a combinatorial library of at least 220 dibenzo[*b,e*][1,4]diazepine[*a,d*]cycloheptene compounds that can be formed by reacting a compound of Formula VII,



with a compound of Formula VIII and



a compound of Formula XII,



wherein X is a halogen; R<sub>1</sub> is selected from the group consisting of hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub> -alkoxyalkyl, and optionally substituted aryl and arylalkyl; R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, and R<sub>5</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>, or R<sub>2</sub> and R<sub>3</sub>, or R<sub>3</sub> and R<sub>4</sub>, or R<sub>4</sub> and R<sub>5</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety; R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, and R<sub>9</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>, or R<sub>6</sub> and R<sub>7</sub>, or R<sub>7</sub> and R<sub>8</sub>, or

R<sub>8</sub> and R<sub>9</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety.


[0014] Also disclosed herein is a pharmaceutical composition comprising a physiologically acceptable carrier, diluent, or excipient, or a combination thereof; and a compound of Formula I, II, or XV.

[0015] Also disclosed herein is a method of treating a neuropsychiatric disorder comprising administering to the patient a therapeutically effective amount of a compound of Formula I, II, or XV.

[0016] Also disclosed herein is a method of treating a neuropsychiatric disorder comprising contacting a therapeutically effective amount of a compound of Formula I, II, or XV with the patient.


[0017] Also disclosed herein is a pharmaceutical composition comprising a compound of Formula I, II, or XV and an neuropsychiatric agent. In some embodiments, the neuropsychiatric agent is selected from the group consisting of a selective serotonin reuptake inhibitor, norepinephrine reuptake inhibitor, dopamine agonist, muscarinic receptor antagonist, antipsychotic agent, serotonin 2A antagonist, and inverse serotonin 2A agonist. In some embodiments, the antipsychotic agent is selected from the group consisting of a phenothiazine, phenylbutylpiperadine, debenzapine, benzisoxidil, and salt of lithium. In some embodiments, the phenothiazine is selected from the group consisting of chlorpromazine (Thorazine®), mesoridazine (Serentil®), prochlorperazine (Compazine®), and thioridazine (Mellaril®). In some embodiments, the phenylbutylpiperadines is selected from the group consisting of haloperidol (Haldol®), and pimozide (Orap®). In some embodiments, the debenzapine is selected from the group consisting of clozapine (Clozaril®), loxapine (Loxitane®), olanzapine (Zyprexa®) and quetiapine (Seroquel®). In some embodiments, the benzisoxidil is selected from the group consisting of resperidone (Resperidal®) and ziprasidone (Geodon®). In some embodiments, the salt of lithium is lithium carbonate. In some embodiments, the antipsychotic agent is selected from the group consisting of Clozaril, Compazine, Etrafon, Geodon, Haldol, Inapsine, Loxitane, Mellaril, Moban, Navane, Orap, Permitil, Prolixin, Phenergan, Reglan, Risperdal, Serentil, Seroquel, Stelazine, Taractan, Thorazine, Triavil, Trilafon, and Zyprexa. In some embodiments, the selective serotonin reuptake inhibitor is selected from the group consisting of fluoxetine, fluvoxamine, sertraline, paroxetine, citalopram, escitalopram,

sibutramine, duloxetine, and venlafaxine, and pharmaceutically acceptable salts or prodrugs thereof. In some embodiments, the norepinephrine reuptake inhibitor is selected from the group consisting of thionisoxetine and reboxetine. In some embodiments, the dopamine agonist is selected from the group consisting of sumatriptan, almotriptan, naratriptan, frovatriptan, rizatriptan, zomitriptan, cabergoline, amantadine, lisuride, pergolide, ropinirole, pramipexole, and bromocriptine. In some embodiments, the inverse serotonin 2A agonist is the compound of Formula XIII, or a related analog thereof.



(XIII)

In some embodiments, the serotonin 2A antagonist is the compound of Formula XIV, or a related analog thereof:



(XIV)


**[0018]** Also disclosed herein is a method of treating neuropsychiatric disorder in a patient comprising administering to the patient a therapeutically effective amount of a pharmaceutical composition comprising a compound of Formula I, II, or XV and an neuropsychiatric agent.

**[0019]** Also disclosed herein is a method of treating neuropsychiatric disorder in a patient comprising administering to the patient a therapeutically effective amount of a compound of Formula I, II, or XV and a therapeutically effective amount of a neuropsychiatric agent. In some embodiments, the administering step comprises administering the compound of Formula I, II, or XV and the neuropsychiatric agent nearly


simultaneously. In other embodiments, the administering step comprises administering one of the compound of Formula I, II, or XV and the neuropsychiatric agent first and then administering the other one of the compound of Formula I, II, or XV and the neuropsychiatric agent. In some embodiments, the neuropsychiatric disorder is selected from the group consisting of schizophrenia and related idiopathic psychoses, anxiety, sleep disorders, appetite disorders, affective disorders such as major depression, bipolar disorder, and depression with psychotic features, and Tourette's Syndrome, drug-induced psychoses, psychoses secondary to neurodegenerative disorders such as Alzheimer's or Huntington's Disease.

### Detailed Description of the Preferred Embodiment


[0020] In the first aspect, the present disclosure is related to a compound of Formula I, II, or XV:



(I)




(II)



(XV)

or a pharmaceutically acceptable salt, ester, amide, or prodrug thereof, wherein:

A is selected from the group consisting of



X is nitrogen, CH, or CH<sub>2</sub>;

X' is C or CH, wherein when X' is C, there is a double bond between X and X' and wherein when X' is CH, there is a single bond between X and X';

each Y is separately selected from the group consisting of nitrogen, oxygen, or CH;

each W is separately selected from the group consisting of nitrogen, CH, oxygen, or sulfur;

each n is separately selected from the group consisting of 0, 1, 2, 3, and 4;

[0021] m is selected from the group consisting of 1, 2, and 3;

each R<sub>1</sub> is separately absent or is separately selected from the group consisting of hydrogen, halogen, amine, optionally substituted C<sub>1-20</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-20</sub> alkenyl, optionally substituted C<sub>2-20</sub> alkynyl, optionally substituted C<sub>1-20</sub>-alkoxyalkyl, and optionally substituted aryl and arylalkyl;

L is absent or is selected from the group consisting of -NH(CH<sub>2</sub>)<sub>n</sub>- and -(CH<sub>2</sub>)<sub>n</sub>-;

a, b, c, and d are each independently selected from the group consisting of carbon, nitrogen, oxygen, and sulfur, or each is independently absent,

provided that at least three of a, b, c, or d are present,

provided that at least one of a, b, c, or d is carbon, and

provided that no two adjacent a, b, c, or d are both oxygen or both sulfur;

e, f, g, and h are each independently selected from the group consisting of carbon, nitrogen, oxygen, and sulfur, or each is independently absent,

provided that at least three of e, f, g, or h are present,

provided that at least one of e, f, g, or h is carbon, and

provided that no two adjacent e, f, g, or h are both oxygen or both sulfur;

R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, and R<sub>5</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>,



or R<sub>2</sub> and R<sub>3</sub>, or R<sub>3</sub> and R<sub>4</sub>, or R<sub>4</sub> and R<sub>5</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety;

R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, and R<sub>9</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>,

or R<sub>6</sub> and R<sub>7</sub>, or R<sub>7</sub> and R<sub>8</sub>, or R<sub>8</sub> and R<sub>9</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety;

Z is selected from the group consisting of NR<sub>11</sub>, oxygen, sulfur, and CH<sub>2</sub>;

R<sub>10</sub> is selected from the group consisting of hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl optionally substituted aryl, optionally substituted arylalkyl, and perhaloalkyl;



R<sub>11</sub> is selected from the group consisting of hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, and optionally substituted arylalkyl;

R<sub>12</sub> and R<sub>13</sub> are separately selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>,

or R<sub>12</sub> and R<sub>13</sub>, taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety.

[0022] Bonds represented by a dashed and solid line represents a bond selected from the group consisting of a carbon-carbon single bond and a carbon-carbon double bond. The dashed bond between X and X' in Formulae I, II, and XV indicates that X and X' may be joined by either a single or a double bond.

[0023] In certain embodiments, the compound of Formulae I and XV does not include clozapine or N-desmethylozapine, the structures of which are shown below:



[0024] In certain embodiments, in compounds of Formulae I and XV, Y is nitrogen or CH. In other embodiments, in compounds of Formula II, Y is nitrogen, oxygen or CH.

[0025] The term “pharmaceutically acceptable salt” refers to a formulation of a compound that does not cause significant irritation to an organism to which it is administered and does not abrogate the biological activity and properties of the compound. Pharmaceutical salts can be obtained by reacting a compound of the invention with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like. Pharmaceutical salts can also be obtained by reacting a compound of the invention with a base to form a salt such as an ammonium salt, an alkali metal salt, such as a sodium or a potassium salt, an alkaline earth metal salt, such as a calcium or a magnesium salt, a salt of organic bases such as dicyclohexylamine, N-methyl-D-glucamine, tris(hydroxymethyl)methylamine, and salts with amino acids such as arginine, lysine, and the like.

[0026] The term “ester” refers to a chemical moiety with formula  $-(R)_n-COOR'$ , where R and R' are independently selected from the group consisting of alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon), and where n is 0 or 1.

[0027] An “amide” is a chemical moiety with formula  $-(R)_n-C(O)NHR'$  or  $-(R)_n-NHC(O)R'$ , where R and R' are independently selected from the group consisting of alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon), and where n is 0 or 1. An amide may be an amino acid or a peptide molecule attached to a molecule of the present invention, thereby forming a prodrug.

[0028] Any amine, hydroxy, or carboxyl side chain on the compounds of the present invention can be esterified or amidified. The procedures and specific groups to be used to achieve this end are known to those of skill in the art and can readily be found in reference sources such as Greene and Wuts, *Protective Groups in Organic Synthesis*, 3<sup>rd</sup> Ed., John Wiley & Sons, New York, NY, 1999, which is incorporated herein in its entirety.

[0029] A “prodrug” refers to an agent that is converted into the parent drug *in vivo*. Prodrugs are often useful because, in some situations, they may be easier to administer than the parent drug. They may, for instance, be bioavailable by oral administration whereas the parent is not. The prodrug may also have improved solubility in pharmaceutical compositions over the parent drug. An example, without limitation, of a prodrug would be a compound of the present invention which is administered as an ester (the “prodrug”) to facilitate transmittal across a cell membrane where water solubility is detrimental to mobility but which then is metabolically hydrolyzed to the carboxylic acid, the active entity, once inside the cell where water-solubility is beneficial. A further example of a prodrug might be a short peptide (polyaminoacid) bonded to an acid group where the peptide is metabolized to reveal the active moiety.

[0030] The term “aromatic” refers to an aromatic group which has at least one ring having a conjugated pi electron system and includes both carbocyclic aryl (e.g., phenyl) and heterocyclic aryl groups (e.g., pyridine). The term includes monocyclic or fused-ring polycyclic (i.e., rings which share adjacent pairs of carbon atoms) groups. The term “carbocyclic” refers to a compound which contains one or more covalently closed ring structures, and that the atoms forming the backbone of the ring are all carbon atoms. The term thus distinguishes carbocyclic from heterocyclic rings in which the ring backbone contains at least one atom which is different from carbon. The term “heteroaromatic” refers to an aromatic group which contains at least one heterocyclic ring.

[0031] As used herein, the term “alkyl” refers to an aliphatic hydrocarbon group. The alkyl moiety may be a “saturated alkyl” group, which means that it does not contain any alkene or alkyne moieties. The alkyl moiety may also be an “unsaturated alkyl” moiety, which means that it contains at least one alkene or alkyne moiety. An “alkene” moiety refers to a group consisting of at least two carbon atoms and at least one carbon-carbon double bond, and an “alkyne” moiety refers to a group consisting of at least two carbon atoms and at least one carbon-carbon triple bond. The alkyl moiety, whether saturated or unsaturated, may be branched, straight chain, or cyclic.

[0032] The alkyl group may have 1 to 20 carbon atoms (whenever it appears herein, a numerical range such as "1 to 20" refers to each integer in the given range; *e.g.*, "1 to 20 carbon atoms" means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 20 carbon atoms, although the present definition also covers the occurrence of the term "alkyl" where no numerical range is designated). The alkyl group may also be a medium size alkyl having 1 to 10 carbon atoms. The alkyl group could also be a lower alkyl having 1 to 5 carbon atoms. The alkyl group of the compounds of the invention may be designated as "C<sub>1</sub>-C<sub>4</sub> alkyl" or similar designations. By way of example only, "C<sub>1</sub>-C<sub>4</sub> alkyl" indicates that there are one to four carbon atoms in the alkyl chain, *i.e.*, the alkyl chain is selected from the group consisting of methyl, ethyl, propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and t-butyl.

[0033] The alkyl group may be substituted or unsubstituted. When substituted, the substituent group(s) is(are) one or more group(s) individually and independently selected from cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, alkoxy, aryloxy, mercapto, alkylthio, arylthio, cyano, halo, carbonyl, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, nitro, silyl, trihalomethanesulfonyl, and amino, including mono- and di-substituted amino groups, and the protected derivatives thereof. Typical alkyl groups include, but are in no way limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tertiary butyl, pentyl, hexyl, ethenyl, propenyl, butenyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like. Wherever a substituent is described as being "optionally substituted" that substituent may be substituted with one of the above substituents.

[0034] The substituent "R" appearing by itself and without a number designation refers to a substituent selected from the group consisting of alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon).

[0035] An "O-carboxy" group refers to a RC(=O)O- group, where R is as defined herein.

[0036] A "C-carboxy" group refers to a -C(=O)OR groups where R is as defined herein.

[0037] An "acetyl" group refers to a -C(=O)CH<sub>3</sub>, group.

[0038] A “trihalomethanesulfonyl” group refers to a  $X_3CS(=O)_2-$  group where X is a halogen.

[0039] A “cyano” group refers to a -CN group.

[0040] An “isocyanato” group refers to a -NCO group.

[0041] A “thiocyanato” group refers to a -CNS group.

[0042] An “isothiocyanato” group refers to a -NCS group.

[0043] A “sulfinyl” group refers to a  $-S(=O)-R$  group, with R as defined herein.

[0044] A “S-sulfonamido” group refers to a  $-S(=O)_2NR-$  group, with R as defined herein.

[0045] A “N-sulfonamido” group refers to a  $RS(=O)_2NH-$  group with R as defined herein.

[0046] A “trihalomethanesulfonarnido” group refers to a  $X_3CS(=O)_2NR-$  group with X and R as defined herein.

[0047] An “O-carbamyl” group refers to a  $-OC(=O)-NR-$  group-with R as defined herein.

[0048] An “N-carbamyl” group refers to a  $ROC(=O)NH-$  group, with R as defined herein.

[0049] An “O-thiocarbamyl” group refers to a  $-OC(=S)-NR-$  group with R as defined herein.

[0050] An “N-thiocarbamyl” group refers to an  $ROC(=S)NH-$  group, with R as defined herein.

[0051] A “C-arnido” group refers to a  $-C(=O)-NR_2$  group with R as defined herein.

[0052] An “N-amido” group refers to a  $RC(=O)NH-$  group, with R as defined herein.

[0053] The term “perhaloalkyl” refers to an alkyl group where all of the hydrogen atoms are replaced by halogen atoms.

[0054] The term “acylalkyl” refers to a  $RC(=O)R'-$  group, with R as defined herein, and R' being a diradical alkylene group. Examples of acylalkyl, without limitation, may include  $CH_3C(=O)CH_2-$ ,  $CH_3C(=O)CH_2CH_2-$ ,  $CH_3CH_2C(=O)CH_2CH_2-$ ,  $CH_3C(=O)CH_2CH_2CH_2-$ , and the like.

[0055] Unless otherwise indicated, when a substituent is deemed to be “optionally substituted,” it is meant that the substituent is a group that may be substituted with one or more group(s) individually and independently selected from cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, alkoxy, aryloxy, mercapto, alkylthio, arylthio, cyano, halo, carbonyl, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, nitro, silyl, trihalomethanesulfonyl, and amino, including mono- and di-substituted amino groups, and the protected derivatives thereof. The protecting groups that may form the protective derivatives of the above substituents are known to those of skill in the art and may be found in references such as Greene and Wuts, above.

[0056] In the present context, the term “cycloalkyl” is intended to cover three-, four-, five-, six-, seven-, and eight- or more membered rings comprising carbon atoms only. A cycloalkyl can optionally contain one or more unsaturated bonds situated in such a way, however, that an aromatic pi-electron system does not arise. Some examples of “cycloalkyl” are the carbocycles cyclopropane, cyclobutane, cyclopentane, cyclopentene, cyclopentadiene, cyclohexane, cyclohexene, 1,3-cyclohexadiene, 1,4-cyclohexadiene, cycloheptane, or cycloheptene.

[0057] The term “heterocyclyl” is intended to mean three-, four-, five-, six-, seven-, and eight- or more membered rings wherein carbon atoms together with from 1 to 3 heteroatoms constitute said ring. A heterocyclyl can optionally contain one or more unsaturated bonds situated in such a way, however, that an aromatic pi-electron system does not arise. The heteroatoms are independently selected from oxygen, sulfur, and nitrogen.

[0058] A heterocyclyl can further contain one or more carbonyl or thiocarbonyl functionalities, so as to make the definition include oxo-systems and thio-systems such as lactams, lactones, cyclic imides, cyclic thioimides, cyclic carbamates, and the like.

[0059] Heterocyclyl rings can optionally also be fused to aryl rings, such that the definition includes bicyclic structures. Typically such fused heterocyclyl groups share one bond with an optionally substituted benzene ring. Examples of benzo-fused heterocyclyl groups include, but are not limited to, benzimidazolidinone, tetrahydroquinoline, and methylenedioxybenzene ring structures.

[0060] Some examples of “heterocyclyls” include, but are not limited to, tetrahydrothiopyran, 4*H*-pyran, tetrahydropyran, piperidine, 1,3-dioxin, 1,3-dioxane, 1,4-dioxin, 1,4-dioxane, piperazine, 1,3-oxathiane, 1,4-oxathiin, 1,4-oxathiane, tetrahydro-1,4-thiazine, 2*H*-1,2-oxazine, maleimide, succinimide, barbituric acid, thiobarbituric acid, dioxopiperazine, hydantoin, dihydrouracil, morpholine, trioxane, hexahydro-1,3,5-triazine, tetrahydrothiophene, tetrahydrofuran, pyrroline, pyrrolidine, pyrrolidone, pyrrolidione, pyrazoline, pyrazolidine, imidazoline, imidazolidine, 1,3-dioxole, 1,3-dioxolane, 1,3-dithiole, 1,3-dithiolane, isoxazoline, isoxazolidine, oxazoline, oxazolidine, oxazolidinone, thiazoline, thiazolidine, and 1,3-oxathiolane. Binding to the heterocycle can be at the position of a heteroatom or via a carbon atom of the heterocycle, or, for benzo-fused derivatives, via a carbon of the benzenoid ring.


[0061] In the present context the term “aryl” is intended to mean a carbocyclic aromatic ring or ring system. Moreover, the term “aryl” includes fused ring systems wherein at least two aryl rings, or at least one aryl and at least one C<sub>3-8</sub>-cycloalkyl share at least one chemical bond. Some examples of “aryl” rings include optionally substituted phenyl, naphthalenyl, phenanthrenyl, anthracenyl, tetralinyl, fluorenyl, indenyl, and indanyl. The term “aryl” relates to aromatic, including, for example, benzenoid groups, connected via one of the ring-forming carbon atoms, and optionally carrying one or more substituents selected from heterocyclyl, heteroaryl, halo, hydroxy, amino, cyano, nitro, alkylamido, acyl, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> hydroxyalkyl, C<sub>1-6</sub> aminoalkyl, C<sub>1-6</sub> alkylamino, alkylsulfenyl, alkylsulfenyl, alkylsulfonyl, sulfamoyl, or trifluoromethyl. The aryl group can be substituted at the *para* and/or *meta* positions. In other embodiments, the aryl group can be substituted at the *ortho* position. Representative examples of aryl groups include, but are not limited to, phenyl, 3-halophenyl, 4-halophenyl, 3-hydroxyphenyl, 4-hydroxyphenyl, 3-aminophenyl, 4-aminophenyl, 3-methylphenyl, 4-methylphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-trifluoromethoxyphenyl, 3-cyanophenyl, 4-cyanophenyl, dimethylphenyl, naphthyl, hydroxynaphthyl, hydroxymethylphenyl, trifluoromethylphenyl, alkoxyphenyl, 4-morpholin-4-ylphenyl, 4-pyrrolidin-1-ylphenyl, 4-pyrazolylphenyl, 4-triazolylphenyl, and 4-(2-oxopyrrolidin-1-yl)phenyl.

[0062] In the present context, the term “heteroaryl” is intended to mean a heterocyclic aromatic group where one or more carbon atoms in an aromatic ring have been replaced with one or more heteroatoms selected from the group comprising nitrogen, sulfur, phosphorous, and oxygen.


[0063] Furthermore, in the present context, the term “heteroaryl” comprises fused ring systems wherein at least one aryl ring and at least one heteroaryl ring, at least two heteroaryl rings, at least one heteroaryl ring and at least one heterocyclyl ring, or at least one heteroaryl ring and at least one cycloalkyl ring share at least one chemical bond.

[0064] The term “heteroaryl” is understood to relate to aromatic, C<sub>3-8</sub> cyclic groups further containing one oxygen or sulfur atom or up to four nitrogen atoms, or a combination of one oxygen or sulfur atom with up to two nitrogen atoms, and their substituted as well as benzo- and pyrido-fused derivatives, for example, connected via one of the ring-forming carbon atoms. Heteroaryl groups can carry one or more substituents, selected from halo, hydroxy, amino, cyano, nitro, alkylamido, acyl, C<sub>1-6</sub>-alkoxy, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-hydroxyalkyl, C<sub>1-6</sub>-aminoalkyl, C<sub>1-6</sub>-alkylamino, alkylsulfinyl, alkylsulfonyl, sulfamoyl, or trifluoromethyl. In some embodiments, heteroaryl groups can be five- and six-membered aromatic heterocyclic systems carrying 0, 1, or 2 substituents, which can be the same as or different from one another, selected from the list above. Representative examples of heteroaryl groups include, but are not limited to, unsubstituted and mono- or di-substituted derivatives of furan, benzofuran, thiophene, benzothiophene, pyrrole, pyridine, indole, oxazole, benzoxazole, isoxazole, benzisoxazole, thiazole, benzothiazole, isothiazole, imidazole, benzimidazole, pyrazole, indazole, tetrazole, quionoline, isoquinoline, pyridazine, pyrimidine, purine and pyrazine, furazan, 1,2,3-oxadiazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, triazole, benzotriazole, pteridine, phenoxazole, oxadiazole, benzopyrazole, quinolizine, cinnoline, phthalazine, quinazoline, and quinoxaline. In some embodiments, the substituents are halo, hydroxy, cyano, O-C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkyl, hydroxy-C<sub>1-6</sub>-alkyl, and amino-C<sub>1-6</sub>-alkyl.

[0065] In certain embodiments, disclosed herein is a compound selected from the following structures:










where R<sub>1</sub>-R<sub>9</sub>, W, Y, and Z are as described herein.

[0066] In certain other embodiments, disclosed herein is a compound selected from the following structures:



where R<sub>1</sub>, W, Y, and Z are as described herein.

[0067] In certain embodiments, disclosed herein is a compound having a structure set forth in Formula III or Formula IV.



where R<sub>1</sub>-R<sub>5</sub>, W, X, X', Y, and Z are as described herein.

[0068] In certain embodiments, none of a, b, c, or d is absent, and the ring formed thereby is a six-membered ring. In further embodiments, none of e, f, g, or h is absent, and consequently, the ring formed thereby is a six-membered ring. In some embodiments, a, b, c, and d are carbon, and the ring formed thereby is an optionally substituted phenyl ring. In further embodiments, e, f, g, and h are carbon, which similarly form an optionally substituted phenyl ring.

[0069] In certain embodiments,  $R_2$  may be selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, and optionally substituted  $C_{1-6}$  alkyloxy. In some embodiments, the alkyl may be selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. In other embodiments, the alkyloxy may be selected from the group consisting of methoxy, ethoxy, propoxy, isopropoxy, butoxy, sec-butoxy, and tert-butoxy. In further embodiments, the halogen may be selected from the group consisting of fluoro, chloro, and bromo. In certain embodiments,  $R_2$  may be selected from the group consisting of hydrogen, methyl, methoxy, and chloro.

[0070] In some embodiments,  $R_3$  may be selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, optionally substituted  $C_{1-6}$  alkyloxy, and  $NO_2$ . The alkyl group may be selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl, while the alkoxy may be selected from the group consisting of methoxy, ethoxy, propoxy, isopropoxy, butoxy, sec-butoxy, and tert-butoxy. In further embodiments, the halogen may be selected from the group consisting of chloro, bromo, and iodo. In other embodiments,  $R_3$  may be selected from the group consisting of hydrogen, methyl, methoxy, chloro, bromo, iodo, and  $NO_2$ .

[0071] In certain embodiments,  $R_4$  may be selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, perhaloalkyl,  $SO_2R_{10}$ , and  $NO_2$ . In some embodiments, the alkyl may be selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. In further embodiments, the perhaloalkyl may be perfluoroalkyl, which in some embodiments, may be trifluoromethyl. In other embodiments, the halogen may be selected from the group consisting of fluoro, chloro, and bromo. When  $R_4$  is  $SO_2R_{10}$ , the  $R_{10}$  may be hydrogen or optionally substituted  $C_{1-6}$  alkyl, which alkyl may be selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. In certain embodiments,  $R_4$  may be selected from the group consisting of hydrogen, methyl, fluoro, chloro, bromo, trifluoromethyl,  $SO_2CH_3$ , and  $NO_2$ .

[0072] In some embodiments,  $R_5$  may be selected from the group consisting of hydrogen, halogen, and optionally substituted  $C_{1-6}$  alkyl. The alkyl may be selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl, while the halogen may be selected from the group consisting of fluoro, chloro, and bromo. In certain embodiments,  $R_5$  may be hydrogen or chloro.

[0073] In certain embodiments,  $R_6$  may be hydrogen or optionally substituted  $C_{1-6}$  alkyl. The alkyl may be selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. In some embodiments,  $R_6$  may be hydrogen.

[0074] In certain embodiments,  $R_7$  may be selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, perhaloalkyl, CN,  $SO_2R_{10}$ , and  $NO_2$ . The alkyl may be selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl, while the halogen may be selected from the group consisting of fluoro, chloro, and bromo. In some embodiments, the perhaloalkyl is perfluoroalkyl, which in some embodiments, may be trifluoromethyl. In the embodiments in which  $R_7$  may be  $SO_2R_{10}$ ,  $R_{10}$  may be hydrogen or optionally substituted  $C_{1-6}$  alkyl, which alkyl may be selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. In certain embodiments,  $R_7$  may be selected from the group consisting of hydrogen, methyl, chloro, trifluoromethyl,  $SO_2CH_3$ , CN, and  $NO_2$ .

[0075] In some embodiments,  $R_8$  may be selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, which alkyl may be selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. The halogen may be selected from the group consisting of fluoro, chloro, and bromo. In certain embodiments,  $R_8$  may be selected from the group consisting of hydrogen, chloro, and bromo.


[0076] Embodiments of the present disclosure include those in which  $R_9$  may be selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, and perhaloalkyl. The alkyl may be selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl. The halogen may be selected from the group consisting of fluoro, chloro, and bromo. The perhaloalkyl may be perfluoroalkyl, which in some embodiments may be trifluoromethyl. In some embodiments,  $R_9$  may be selected from the group consisting of hydrogen, chloro, methyl, and trifluoromethyl.

[0077] In some embodiments,  $R_{11}$  may be selected from the group consisting of hydrogen, optionally substituted  $C_{1-6}$  alkyl, and optionally substituted aryl. The alkyl may

be selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl, while the aryl may be phenyl or naphthyl. In other embodiments,  $R_1$  may be a heteroaryl. In certain embodiments,  $R_1$  may be hydrogen. In certain embodiments,  $R_1$  is absent.

[0078] In some embodiments, X may be nitrogen. In other embodiments, Y may be NH and W may be nitrogen or CH.

[0079] In some embodiments of the compounds of Formula I or Formula XV, L is absent or is selected from the group consisting of  $-\text{NHCH}_2-$ ,  $-\text{NH}-$ , and  $-\text{CH}_2-$ . In some embodiments of the compounds of Formula I or Formula XV, A is selected from the group consisting of:



where n is selected from the group consisting of 0, 1, and 2.

[0080] Some embodiments of the compounds of Formula I, Formula II, or Formula XV, include:

2,7-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,  
 2-Chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,  
 2,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,  
 8-Bromo-2-chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,  
 2-Chloro-11-(piperazin-1-yl)-8-trifluoromethyl-5H-dibenzo[b,e][1,4]diazepine,  
 6-Chloro-11-(piperazin-1-yl)-8-trifluoromethyl-5H-dibenzo[b,e][1,4]diazepine,  
 7-Chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,  
 8-Bromo-1-chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,  
 8-Bromo-2-methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,  
 4,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,  
 8-Chloro-2-methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,  
 8-Chloro-2-fluoro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,  
 3,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,  
 2-Bromo-8-chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,  
 3,7-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine,



8-Bromo-3-chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 3-Chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 3-Chloro-11-(piperazin-1-yl)-8-trifluoromethyl-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 7-Chloro-2-methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 2-Methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 2-Methyl-11-(piperazin-1-yl)-8-trifluoromethyl-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-4-methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 1,8-Dichloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Bromo-5-methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 7,8-Dichloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 11-(Piperazin-1-yl)-8-trifluoromethyl-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 11-(Piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Fluoro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 11-(Piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine-8-carbonitrile,  
 8-Bromo-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 3-Fluoro-6-piperazin-1-yl-11*H*-dibenzo[*b,e*]azepine,  
 2-(Trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5*H*-  
 dibenzo[*b,e*][1,4]diazepine,  
 2-(Trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5*H*-  
 dibenzo[*b,e*][1,4]oxazepine,  
 8-Chloro-2-(trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5*H*-  
 dibenzo[*b,e*][1,4]diazepine,  
 8-(Trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5*H*-  
 dibenzo[*b,e*][1,4]diazepine,  
 11-(Piperazin-1-yl)-dibenzo[*b,f*][1,4]thiazepin,  
 11-(Piperazin-1-yl)-2,3-dihydro-1,4-benzodioxino[6,7-*b*][1,4]benzothiazepin,  
 8-Chloro-11-[1,4]diazepam-1-yl-5*H*-dibenzo[*b,e*][1,4]diazepine,  
*N'*-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)-*N,N*-dimethyl-ethane-1,2-  
 diamine,  
*N'*-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)-*N,N*-diethyl-ethane-1,2-  
 diamine,  
 8-Chloro-11-(4-methyl-[1,4]diazepam-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,

8-Chloro-2-methoxy-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
*N*<sup>7</sup>-(5*H*-Dibenzo[*b,e*][1,4]diazepine-11-yl)-*N,N*-dimethyl-ethane-1,2-diamine,  
 11-[1,4]Diazepam-1-yl-5*H*-dibenzo[*b,e*][1,4]diazepine,  
*N*<sup>7</sup>-(8-Fluoro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)-*N,N*-dimethyl-ethane-1,2-diamine,  
 8-Fluoro-11-[1,4]diazepam-1-yl-5*H*-dibenzo[*b,e*][1,4]diazepine,  
*N*<sup>7</sup>-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)-*N*-methyl-ethane-1,2-diamine,  
 8-Chloro-11-(trans-2,5-dimethyl-piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-(3,5-dimethyl-piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-(3-methyl-piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-(3-phenyl-piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-5-methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-5-benzyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Iodo-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 2-Iodo-8-chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Phenyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-(piperidin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-(morpholin-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 5-Allyl-8-chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 6-Chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-5-piperazin-1-yl-11*H*-benzo[*b*]pyrido[2,3-*e*][1,4]diazepine,  
 2-Chloro-10-piperazin-1-yl-5*H*-dibenzo[*b,f*]azepin,  
 8-Chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]thiazepine,  
 8-Chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Chloro-11-(4-methyl-piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 3-Chloro-6-piperazin-1-yl-11*H*-dibenzo[*b,e*]azepine,  
 8-Bromo-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 11-(Piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 7-Chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Chloro-3-methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Bromo-3-methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 3-Methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 7-Chloro-3-methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,

8-Chloro-4-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Bromo-4-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 4-Methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 2-Bromo-8-chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 2,8-Dibromo-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 2-Bromo-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 2-Bromo-7-chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 11-(Piperazin-1-yl)-8-trifluoromethyl-dibenzo[*b,f*][1,4]oxazepine,  
 4-Methyl-11-(piperazin-1-yl)-8-trifluoromethyl-dibenzo[*b,f*][1,4]oxazepine,  
 8-Fluoro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Fluoro-3-methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Fluoro-4-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 2-Bromo-8-fluoro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 3-Methoxy-8-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 4,8-Dimethyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 3-Methoxy-11-(piperazin-1-yl)-8-trifluoromethyl-dibenzo[*b,f*][1,4]oxazepine,  
 2-Bromo-11-(piperazin-1-yl)-8-trifluoromethyl-dibenzo[*b,f*][1,4]oxazepine,  
 6-Chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 2-Bromo-8-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 7-Chloro-4-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Phenyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Chloro-11-(piperidin-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine  
 5-Benzyl-8-chloro-11-(piperidin-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Bromo-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one,  
 5,10-Dihydro-dibenzo[*b,e*][1,4]diazepine-11-one,  
 8-Fluoro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one,  
 8,5-Dichloro-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-methylsulfanyl-5*H*-dibenzo[*b,e*][1,4]diazepine  
 (8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-(S)-1-pyrrolidin-2-yl-methyl-amine,  
 1-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-piperidine-4-yl-amine,  
 1-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-pyrrolidin-3-yl-amine,  
 (8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-(R)-1-pyrrolidin-2-yl-methyl-amine,


(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-pyrrolidin-3-yl-amine,  
 8-Chloro-11-(2,5-diaza-bicyclo[2.2.1]hept-2-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 Acetidin-3-yl-(8-chloro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)amine,  
 7-Bromo-4-(piperazin-1-yl)-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepine,  
 7-Bromo-2-methyl-(piperazin-1-yl)-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepine  
 7-Bromo-2-phenyl-4-(piperazine-1-yl)-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepine,  
 7-Bromo-10-(piperazin-1-yl)-1,2,3,3a,4,10a-hexahydro-  
 benzo[*b*]cyclopenta[*e*][1,4]diazepine,  
 8-Chloro-11-(4-fluorobenzyl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-(4-fluorophenyl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-(4-nonylphenyl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-(pyridin-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine, and  
 8-Chloro-11-(1*H*-pyrazol-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine.

[0081] In another aspect, the present disclosure is directed to a method of synthesizing a compound of Formula V or Formula VI,




comprising

reacting a compound of Formula VII




with a compound of Formula VIII




to form a fused ring compound of Formula IX,





and reacting the compound of Formula IX with a compound of Formula X




to obtain a compound of Formula V or VI;


wherein X is a halogen; and R<sub>1</sub>-R<sub>9</sub> are as defined herein. In some embodiments, the compound of Formula V synthesized according to the disclosed method is clozapine while in other embodiments, the compound is N-desmethylozapine. In certain other embodiments, the compound of Formula V synthesized according to the disclosed method does not include clozapine or N-desmethylozapine.

**[0082]** Consistent with this aspect, Schemes 1 and 2 depict the synthesis of some of the compounds disclosed herein. The first series of steps generating the intermediate lactam have been described by, *inter alia*, Liao *et al. J. Med. Chem.* **1997**, *40*, 4146-4153. The last step has been described by e.g. Liao *et al. J. Med. Chem.* **1999**, *42*, 2235-2244. Both of these references are hereby incorporated herein by reference in their entirety, including any drawings.

Scheme 1



Scheme 2




[0083] In certain embodiments of the invention the building blocks A and B are selected from but not limited to


A


B



[0084] In another aspect, the present disclosure relates to a combinatorial library of at least 10, or at least 30, or at least 50, or at least 100, or at least 200, or at least 220 dibenzo[*b,e*][1,4]diazepine compounds that can be formed by reacting a compound of Formula VII,



with a compound of Formula VIII and




a compound of Formula XI,




wherein X is a halogen; W is nitrogen, CH, oxygen, or sulfur; n is 1, 2, 3, or 4 and R<sub>1</sub>-R<sub>9</sub> are as defined herein. In some embodiments, the combinatorial library includes clozapine and/or N-desmethylozapine. In certain other embodiments, the combinatorial library does not include clozapine or N-desmethylozapine.


[0085] In another aspect, the present disclosure relates to a combinatorial library of at least 10, or at least 30, or at least 50, or at least 100, or at least 200, or at least 220 dibenzo[*b,e*][1,4]diazepine compounds that can be formed by reacting a compound of Formula VII,



with a compound of Formula VIII and



a compound of Formula XII,



wherein X is a halogen; W is nitrogen, CH, oxygen, or sulfur; n is 1, 2, 3, or 4; and R<sub>1</sub>-R<sub>9</sub> are as defined herein.

[0086] As used herein, a “combinatorial library” refers to all the compounds formed by the reaction of each compound of one dimension with a compound in each of the other dimensions in a multi-dimensional array of compounds. In the context of the present disclosure, the array is three dimensional, with one dimension representing all the compounds of Formula VII, the second dimension representing all the compounds of Formula VIII, and the third dimension representing all the compounds of Formula X. Each compound of Formula VII may be reacted with each and every compound of Formula VIII and each and every compound of Formula X in order to form a substituted compound of Formula V or VI. All compounds falling within the scope of Formula V or VI formed in this way are within the scope of the present disclosure. Also within the scope of the present disclosure are smaller combinatorial libraries formed by the reaction of some of, or all of, the compounds of Formula VII with some of, or all of, the compounds of Formula VIII and some of, or all of, the compounds of Formula X.

[0087] In some embodiments, compounds of Formulae I, II, or XV, as disclosed and described herein, may be capable of modulating the activity of a muscarinic receptor.

[0088] The term “modulate” refers to the ability of a compound disclosed herein to alter the function of a muscarinic receptor. A modulator may activate the activity of a muscarinic receptor, may activate or inhibit the activity of a muscarinic receptor depending on the concentration of the compound exposed to the muscarinic receptor, or may inhibit the activity of a muscarinic receptor. The term “modulate” also refers to altering the function of a muscarinic receptor by increasing or decreasing the probability that a complex forms between a muscarinic receptor and a natural binding partner. A modulator may increase the probability that such a complex forms between the muscarinic receptor and the natural binding partner, may increase or decrease the probability that a complex forms between the muscarinic receptor and the natural binding partner depending on the concentration of the compound exposed to the muscarinic receptor, and or may decrease the probability that a complex forms between the muscarinic receptor and the natural binding partner. In some embodiments, modulation of the muscarinic receptor may be assessed using Receptor Selection and Amplification Technology (R-SAT) as described in U.S. Patent No. 5,707,798, the disclosure of which is incorporated herein by reference in its entirety.

[0089] The term “activate” refers to increasing the cellular function of a muscarinic receptor. The term “inhibit” refers to decreasing the cellular function of a

muscarinic receptor. The muscarinic receptor function may be the interaction with a natural binding partner or catalytic activity.

[0090] The term “contacting” as used herein refers to bringing a compound disclosed herein and a target muscarinic receptor together in such a manner that the compound can affect the activity of the muscarinic receptor, either directly; i.e., by interacting with the muscarinic receptor itself, or indirectly; i.e., by interacting with another molecule on which the activity of the muscarinic receptor is dependent. Such “contacting” can be accomplished in a test tube, a petri dish or the like. In a test tube, contacting may involve only a compound and a muscarinic receptor of interest or it may involve whole cells. Cells may also be maintained or grown in cell culture dishes and contacted with a compound in that environment. In this context, the ability of a particular compound to affect a muscarinic receptor related disorder; i.e., the  $IC_{50}$  of the compound can be determined before use of the compounds in vivo with more complex living organisms is attempted. For cells outside the organism, multiple methods exist, and are well-known to those skilled in the art, to get the muscarinic receptors in contact with the compounds including, but not limited to, direct cell microinjection and numerous transmembrane carrier techniques. The term “contacting” can also refer to bringing a compound disclosed herein to contact with a target muscarinic receptor in vivo. Thus, if a compound disclosed herein, or a prodrug thereof, is administered to an organism and the compound is brought together with a muscarinic receptor within the organism, such contacting is within the scope of the present disclosure.

[0091] In some embodiments, the compound of Formulae I, II, or XV may be an agonist of said receptor, while in other embodiments, the compound may be an antagonist of said receptor. In yet other embodiments, the compound may be a partial agonist of said receptor. A compound that is a partial agonists may in some cases be a partial activator of a receptor, while in other cases may be a partial repressor of a receptor. In yet other circumstances, the compound may be a tissue-specific modulator, while in other circumstances, the compound may be a gene-specific modulator.

[0092] Certain of the compounds disclosed herein may exist as stereoisomers including optical isomers. The scope of the present disclosure includes all stereoisomers and both the racemic mixtures of such stereoisomers as well as the individual enantiomers that may be separated according to methods that are well known to those of ordinary skill in the art.

[0093] In another aspect, the present disclosure relates to a pharmaceutical composition comprising a physiologically acceptable carrier, diluent, or excipient, or a combination thereof; and a compound of Formulae I, II, or XV.

[0094] The term “pharmaceutical composition” refers to a mixture of a compound of the invention with other chemical components, such as diluents or carriers. The pharmaceutical composition facilitates administration of the compound to an organism. Multiple techniques of administering a compound exist in the art including, but not limited to, oral, injection, aerosol, parenteral, and topical administration. Pharmaceutical compositions can also be obtained by reacting compounds with inorganic or organic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

[0095] The term “carrier” defines a chemical compound that facilitates the incorporation of a compound into cells or tissues. For example dimethyl sulfoxide (DMSO) is a commonly utilized carrier as it facilitates the uptake of many organic compounds into the cells or tissues of an organism.

[0096] The term “diluent” defines chemical compounds diluted in water that will dissolve the compound of interest as well as stabilize the biologically active form of the compound. Salts dissolved in buffered solutions are utilized as diluents in the art. One commonly used buffered solution is phosphate buffered saline because it mimics the salt conditions of human blood. Since buffer salts can control the pH of a solution at low concentrations, a buffered diluent rarely modifies the biological activity of a compound.

[0097] The term “physiologically acceptable” defines a carrier or diluent that does not abrogate the biological activity and properties of the compound.

[0098] The pharmaceutical compositions described herein can be administered to a human patient *per se*, or in pharmaceutical compositions where they are mixed with other active ingredients, as in combination therapy, or suitable carriers or excipient(s). Techniques for formulation and administration of the compounds of the instant application may be found in “Remington’s Pharmaceutical Sciences,” Mack Publishing Co., Easton, PA, 18th edition, 1990.

[0099] Suitable routes of administration may, for example, include oral, rectal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular,

subcutaneous, intravenous, intramedullary injections, as well as intrathecal, direct intraventricular, intraperitoneal, intranasal, or intraocular injections.

[0100] Alternately, one may administer the compound in a local rather than systemic manner, for example, via injection of the compound directly in the renal or cardiac area, often in a depot or sustained release formulation. Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome coated with a tissue-specific antibody. The liposomes will be targeted to and taken up selectively by the organ.

[0101] The pharmaceutical compositions of the present invention may be manufactured in a manner that is itself known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or tableting processes.

[0102] Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. Any of the well-known techniques, carriers, and excipients may be used as suitable and as understood in the art; *e.g.*, in Remington's Pharmaceutical Sciences, above.

[0103] For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

[0104] For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by mixing one or more solid excipient with pharmaceutical combination of the invention, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch,



potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

[0105] Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

[0106] Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

[0107] For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

[0108] For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, *e.g.*, gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[0109] The compounds may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or

emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

[0110] Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

[0111] Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

[0112] The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides.

[0113] In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[0114] A pharmaceutical carrier for the hydrophobic compounds of the invention is a cosolvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. A common cosolvent system used is the VPD co-solvent system, which is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant Polysorbate 80™, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used instead of POLYSORBATE 80™; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may

replace polyethylene glycol, *e.g.*, polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose.

[0115] Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

[0116] Many of the compounds used in the pharmaceutical combinations of the invention may be provided as salts with pharmaceutically compatible counterions. Pharmaceutically compatible salts may be formed with many acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, *etc.* Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free acid or base forms.

[0117] Pharmaceutical compositions suitable for use in the present invention include compositions where the active ingredients are contained in an amount effective to achieve its intended purpose. More specifically, a therapeutically effective amount means an amount of compound effective to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the subject being treated. Determination of a therapeutically effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

[0118] The exact formulation, route of administration and dosage for the pharmaceutical compositions of the present invention can be chosen by the individual physician in view of the patient's condition. (See *e.g.*, Fingl *et al.* 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p. 1). Typically, the dose range of the composition administered to the patient can be from about 0.5 to 1000 mg/kg of the patient's body weight. The dosage may be a single one or a series of two or more given in the course of one or more days, as is needed by the patient. Note that for almost all of the

specific compounds mentioned in the present disclosure, human dosages for treatment of at least some condition have been established. Thus, in most instances, the present invention will use those same dosages, or dosages that are between about 0.1% and 500%, more preferably between about 25% and 250% of the established human dosage. Where no human dosage is established, as will be the case for newly-discovered pharmaceutical compounds, a suitable human dosage can be inferred from ED<sub>50</sub> or ID<sub>50</sub> values, or other appropriate values derived from *in vitro* or *in vivo* studies, as qualified by toxicity studies and efficacy studies in animals.

[0119] Although the exact dosage will be determined on a drug-by-drug basis, in most cases, some generalizations regarding the dosage can be made. The daily dosage regimen for an adult human patient may be, for example, an oral dose of between 0.1 mg and 500 mg of each ingredient, preferably between 1 mg and 250 mg, e.g. 5 to 200 mg or an intravenous, subcutaneous, or intramuscular dose of each ingredient between 0.01 mg and 100 mg, preferably between 0.1 mg and 60 mg, e.g. 1 to 40 mg of each ingredient of the pharmaceutical compositions of the present invention or a pharmaceutically acceptable salt thereof calculated as the free base, the composition being administered 1 to 4 times per day. Alternatively the compositions of the invention may be administered by continuous intravenous infusion, preferably at a dose of each ingredient up to 400 mg per day. Thus, the total daily dosage by oral administration of each ingredient will typically be in the range 1 to 2000 mg and the total daily dosage by parenteral administration will typically be in the range 0.1 to 400 mg. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more, or for months or years.

[0120] Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the modulating effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations.

[0121] Dosage intervals can also be determined using MEC value. Compositions should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%.

[0122] In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

[0123] The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

[0124] The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser may also be accompanied with a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the drug for human or veterinary administration. Such notice, for example, may be the labeling approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

[0125] In another aspect, the present disclosure is related to a method of treating a neuropsychiatric disorder comprising administering to said patient a therapeutically effective amount of a compound of Formula I or II. In some embodiments, the neuropsychiatric disorder is selected from the group consisting of schizophrenia and related idiopathic psychoses, anxiety, sleep disorders, appetite disorders, affective disorders such as major depression, bipolar disorder, and depression with psychotic features, and Tourette's Syndrome, drug-induced psychoses, psychoses secondary to neurodegenerative disorders such as Alzheimer's or Huntington's Disease.

[0126] In some embodiments, the compound of Formulae I or XV is clozapine while in other embodiments, the compound is N-desmethylozapine. In certain other embodiments, the compound of Formulae I or XV does not include clozapine or N-desmethylozapine.

[0127] In yet another aspect, the present disclosure is related to a method of treating a neuropsychiatric disorder comprising contacting a therapeutically effective amount of a compound of Formulae I, II, or XV with said patient .

**[0128]** In some embodiments, the present disclosure relates to a pharmaceutical composition comprising a compound of Formula I, II, or XV and a neuropsychiatric agent. As used herein, a “neuropsychiatric agent” refers to a compound, or a combination of compounds, that affects the neurons in the brain either directly or indirectly, or affects the signal transmitted to the neurons in the brain. Neuropsychiatric agents, therefore, may affect a person’s psyche, such as the person’s mood, perception, nociception, cognition, alertness, memory, etc. In certain embodiments, the neuropsychiatric agent may be selected from the group consisting of a selective serotonin reuptake inhibitor, norepinephrine reuptake inhibitor, dopamine agonist, antipsychotic agent, serotonin 2A antagonists, and inverse serotonin 2A agonists.


**[0129]** In some embodiments, the antipsychotic agent may be selected from the group consisting of a phenothiazine, phenylbutylpiperadine, debenzapine, benzisoxidil, and salt of lithium. The phenothiazine group of compounds may be selected from the group consisting of chlorpromazine (Thorazine®), mesoridazine (Serentil®), prochlorperazine (Compazine®), and thioridazine (Mellaril®). The phenylbutylpiperadine group of compounds may be selected from the group consisting of haloperidol (Haldol®), and pimozide (Orap®). The debenzapine group of compounds may be selected from the group consisting of clozapine (Clozaril®), loxapine (Loxitane®), olanzapine (Zyprexa®) and quetiapine (Seroquel®). The benzisoxidil group of compounds may be selected from the group consisting of risperidone (Risperidal®) and ziprasidone (Geodon®). The salt of lithium may be lithium carbonate. In some embodiments, the antipsychotic agent may be selected from the group consisting of Aripiprazole (Abilify), Clozapine, Clozaril, Compazine, Etrafon, Geodon, Haldol, Inapsine, Loxitane, Mellaril, Moban, Navane, Olanzapine (Zyprexa), Orap, Permitil, Prolixin, Phenergan, Quetiapine (Seroquel), Reglan, Risperdal, Serentil, Seroquel, Stelazine, Taractan, Thorazine, Triavil, Trilafon, and Zyprexa, or pharmaceutically acceptable salts thereof.

**[0130]** In certain embodiments, the selective serotonin reuptake inhibitor is selected from the group consisting of fluoxetine, fluvoxamine, sertraline, paroxetine, citalopram, escitalopram, sibutramine, duloxetine, and venlafaxine, and pharmaceutically acceptable salts or prodrugs thereof.


**[0131]** In other embodiments, the norepinephrine reuptake inhibitor is selected from the group consisting of thionisoxetine and reboxetine.

[0132] In further embodiments, the dopamine agonist is selected from the group consisting of sumatriptan, almotriptan, naratriptan, frovatriptan, rizatriptan, zomitriptan, cabergoline, amantadine, lisuride, pergolide, ropinirole, pramipexole, and bromocriptine.

[0133] In another embodiment, the inverse serotonin 2A agonist is ACP-103 or an analog thereof. By "ACP-103," it is meant the compound of Formula XIII.



[0134] In another embodiment, the serotonin 2A antagonist is M 100,907 or an analog thereof. By "M 100,907," it is meant the compound of Formula XIV.



[0135] In another aspect, the present disclosure is directed to a method of treating neuropsychiatric disorder in a patient comprising administering to said patient a therapeutically effective amount of a pharmaceutical composition comprising a compound of Formula I, II, or XV and a neuropsychiatric agent. In yet another aspect, the present disclosure is directed to a method of treating neuropsychiatric disorder in a patient comprising administering to said patient a therapeutically effective amount of a compound of Formula I, II, or XV and a therapeutically effective amount of a neuropsychiatric agent.

[0136] In certain embodiments, the patient may be a mammal. The mammal may be selected from the group consisting of mice, rats, rabbits, guinea pigs, dogs, cats, sheep, goats, cows, primates, such as monkeys, chimpanzees, and apes, and humans. In some embodiments, the patient is a human.

[0137] In some embodiments, the administering step in the above methods comprises administering said compound of Formula I, II, or XV and said neuropsychiatric

agent nearly simultaneously. These embodiments include those in which the compound of Formula I or II and the neuropsychiatric agent are in the same administrable composition, i.e., a single tablet, pill, or capsule, or a single solution for intravenous injection, or a single drinkable solution, or a single dragee formulation or patch, contains both compounds. The embodiments also include those in which each compound is in a separate administrable composition, but the patient is directed to take the separate compositions nearly simultaneously, i.e., one pill is taken right after the other or that one injection of one compound is made right after the injection of another compound, etc.

[0138] In other embodiments the administering step comprises administering one of the compound of Formula I, II, or XV and the neuropsychiatric agent first and then administering the other one of the compound of Formula I, II, or XV and the neuropsychiatric agent. In these embodiments, the patient may be administered a composition comprising one of the compounds and then at some time, a few minutes or a few hours, later be administered another composition comprising the other one of the compounds. Also included in these embodiments are those in which the patient is administered a composition comprising one of the compounds on a routine or continuous basis while receiving a composition comprising the other compound occasionally.

[0139] In certain embodiments, the neuropsychiatric disorder to be treated by the methods and the compounds of the present disclosure is selected from the group consisting of schizophrenia and related idiopathic psychoses, anxiety, sleep disorders, appetite disorders, affective disorders such as major depression, bipolar disorder, and depression with psychotic features, and Tourette's Syndrome, drug-induced psychoses, psychoses secondary to neurodegenerative disorders such as Alzheimer's or Huntington's Disease.

## EXAMPLES

### Example 1 – General procedure 1 (GP1)

[0140] A mixture of an aminobenzoic acid (1 eq.), a 2-fluoronitrobenzene (3 eq.) and  $\text{Cs}_2\text{CO}_3$  (3 eq.) in DMF was heated to 140°C for 1 hour, and then allowed to obtain room temperature. The mixture was diluted with water and washed with EtOAc (2 x).


[0141] EtOH and  $\text{Na}_2\text{S}_2\text{O}_4$  (5 eq.) was added to the aqueous phase and the resulting mixture was stirred for 1 h. Aqueous HCl (2 M) was added to the mixture and then the aqueous phase was extracted with EtOAc (3 x) and the combined organic phases were concentrated.



[0142] The residue was taken up in  $\text{CH}_2\text{Cl}_2$  and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (3 eq.) was added and the resulting mixture was stirred at room temperature for 1 h, and then concentrated. The residue was diluted with EtOAc, washed with aqueous NaOH (2 M) and concentrated.


[0143] The residue was taken up in dioxane and added to a mixture of  $\text{TiCl}_4$  (1.1 eq., 1 M in toluene) and piperazine (5 eq.) in dioxane at  $50^\circ\text{C}$ . The resulting mixture was stirred at  $100^\circ\text{C}$  over night, and then allowed to obtain room temperature. Aqueous HCl (2 M) was added to the mixture until the solution became acidic and then the aqueous phase was extracted with EtOAc (2 x). Aqueous NaOH (2 M) was added to the aqueous phase until a basic solution was obtained and the resulting suspension was extracted with EtOAc (3 x). The combined organic phases were concentrated and purified by HPLC.

Example 2 – 2,7-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO85F1)




[0144] 4-Chloro-2-fluoronitrobenzene (263 mg, 1.5 mmol) and 2-amino-5-chlorobenzoic acid (86 mg, 0.5 mmol) were reacted according to GP1 to give 6.1 mg of the title compound (166 JO85F1). MS (ESI) 347 ( $\text{MH}^+$ ). Purity for  $\text{MH}^+$  (UV/MS) 100/85.

Example 3 – 2-Chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO85F6)




[0145] 2-Fluoronitrobenzene (212 mg, 1.5 mmol) and 2-amino-5-chlorobenzoic acid (86 mg, 0.5 mmol) were reacted according to GP1 to give 5.3 mg of the title compound (166 JO85F6). MS (ESI) 313 ( $\text{MH}^+$ ). Purity for  $\text{MH}^+$  (UV/MS) 100/95.

Example 4 – 2,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO85F2)




[0146] 5-Chloro-2-fluoronitrobenzene (263 mg, 1.5 mmol) and 2-amino-5-chlorobenzoic acid (86 mg, 0.5 mmol) were reacted according to GP1 to give 4.8 mg of the title compound (166 JO85F2). MS (ESI) 347 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/99.

Example 5 – 8-Bromo-2-chloro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (166JO85F3)




[0147] 5-Bromo-2-fluoronitrobenzene (330 mg, 1.5 mmol) and 2-amino-5-chlorobenzoic acid (86 mg, 0.5 mmol) were reacted according to GP1 to give 8.0 mg of the title compound (166 JO85F3). MS (ESI) 391 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/96.

Example 6 – 2-Chloro-11-(piperazin-1-yl)-8-trifluoromethyl-5H-dibenzo[*b,e*][1,4]diazepine (166JO85F7)




[0148] 4-Fluoro-3-nitrobenzotrifluoride (314 mg, 1.5 mmol) and 2-amino-5-chlorobenzoic acid (86 mg, 0.5 mmol) were reacted according to GP1 to give 0.3 mg of the title compound (166 JO85F7). MS (ESI) 381 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/95.

Example 7 – 6-Chloro-11-(piperazin-1-yl)-8-trifluoromethyl-5H-dibenzo[*b,e*][1,4]diazepine (189JO77B)




[0149] 3-Chloro-4-fluoro-5-nitrobenzotrifluoride (366 mg, 1.5 mmol) and 2-aminobenzoic acid (69 mg, 0.5 mmol) were reacted according to GP1 to give 28 mg of the title compound (189JO77B). MS (ESI) 381 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/100.

Example 8 – 7-Chloro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE35B)




[0150] 4-Chloro-2-fluoronitrobenzene (528 mg, 3.0 mmol) and 2-aminobenzoic acid (138 mg, 1.0 mmol) were reacted according to GP1 to give 5.0 mg of the title compound (160FE35B). MS (ESI) 313 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/86.

Example 9 – 8-Bromo-1-chloro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE36A)




[0151] 5-Bromo-2-fluoronitrobenzene (660 mg, 3.0 mmol) and 2-amino-6-chlorobenzoic acid (172 mg, 1.0 mmol) were reacted according to GP1 to give 5.0 mg of the title compound (160FE36A). MS (ESI) 391 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 94/87.

Example 10 – 8-Bromo-2-methyl-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE40C)




[0152] 5-Bromo-2-fluoronitrobenzene (660 mg, 3.0 mmol) and 2-amino-5-methylbenzoic acid (152 mg, 1.0 mmol) were reacted according to GP1 to give 7.9 mg of the title compound (160FE40C). MS (ESI) 371 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 11 – 4,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE41A)




[0153] 5-Chloro-2-fluoronitrobenzene (527 mg, 3.0 mmol) and 2-amino-3-chlorobenzoic acid (172 mg, 1.0 mmol) were reacted according to GP1 to give 4.6 mg of the title compound (160FE41A). MS (ESI) 347 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 95/70.

Example 12 – 8-Chloro-2-methyl-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE41B)




[0154] 5-Chloro-2-fluoronitrobenzene (527 mg, 3.0 mmol) and 2-amino-5-methylbenzoic acid (151 mg, 1.0 mmol) were reacted according to GP1 to give 7.1 mg of the title compound (160FE41B). MS (ESI) 327 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/94.

Example 13 – 8-Chloro-2-fluoro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE42A-F3)




[0155] 5-Chloro-2-fluoronitrobenzene (264 mg, 1.5 mmol) and 2-amino-5-fluorobenzoic acid (78 mg, 0.5 mmol) were reacted according to GP1 to give 21 mg of the title compound (160FE42A-F3). MS (ESI) 331 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/98.

Example 14 – 3,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE42B-F4)




[0156] 5-Chloro-2-fluoronitrobenzene (264 mg, 1.5 mmol) and 2-amino-4-chlorobenzoic acid (86 mg, 0.5 mmol) were reacted according to GP1 to give 9.4 mg of the title compound (160FE42B-F4). MS (ESI) 347 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/97.

Example 15 – 2-Bromo-8-chloro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE43A-F6)




[0157] 5-Chloro-2-fluoronitrobenzene (528 mg, 3.0 mmol) and 2-amino-5-bromobenzoic acid (216 mg, 1.0 mmol) were reacted according to GP1 to give 20 mg of the title compound (160FE43A-F6). MS (ESI) 391 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 16 – 3,7-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE58D1)




[0158] 4-Chloro-2-fluoronitrobenzene (263 mg, 1.5 mmol) and 2-amino-4-chlorobenzoic acid (86 mg, 0.5 mmol) were reacted according to GP1 to give 3.1 mg of the title compound (160FE58D1). MS (ESI) 347 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 63/83.

Example 17 – 8-Bromo-3-chloro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE58D3)




[0159] 5-Bromo-2-fluoronitrobenzene (330 mg, 1.5 mmol) and 2-amino-4-chlorobenzoic acid (86 mg, 0.5 mmol) were reacted according to GP1 to give 1.1 mg of the title compound (160FE58D3). MS (ESI) 391 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 90/85.

Example 18 – 3-Chloro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE58D6)




[0160] 2-Fluoronitrobenzene (212 mg, 1.5 mmol) and 2-amino-4-chlorobenzoic acid (86 mg, 0.5 mmol) were reacted according to GP1 to give 2.2 mg of the title compound (160FE58D6). MS (ESI) 313 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 90/100.

Example 19 – 3-Chloro-11-(piperazin-1-yl)-8-trifluoromethyl-5H-dibenzo[b,e][1,4]diazepine (160FE58D7)




[0161] 4-Fluoro-3-nitrobenzotrifluoride (314 mg, 1.5 mmol) and 2-amino-4-chlorobenzoic acid (86 mg, 0.5 mmol) were reacted according to GP1 to give 2.0 mg of the title compound (160FE58D7). MS (ESI) 381 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 20 – 7-Chloro-2-methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE58E1)




[0162] 4-Chloro-2-fluoronitrobenzene (263 mg, 1.5 mmol) and 2-amino-5-methylbenzoic acid (76 mg, 0.5 mmol) were reacted according to GP1 to give 1.1 mg of the title compound (160FE58E1). MS (ESI) 327 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/90.

Example 21 – 2-Methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE58E6)




[0163] 4-Fluoronitrobenzene (212 mg, 1.5 mmol) and 2-amino-5-methylbenzoic acid (76 mg, 0.5 mmol) were reacted according to GP1 to give 6.8 mg of the title compound (160FE58E6). MS (ESI) 293 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 22 – 2-Methyl-11-(piperazin-1-yl)-8-trifluoromethyl-5H-dibenzo[*b,e*][1,4]diazepine (160FE58E7)




[0164] 4-Fluoro-3-nitrobenzotrifluoride (314 mg, 1.5 mmol) and 2-amino-5-methylbenzoic acid (76 mg, 0.5 mmol) were reacted according to GP1 to give 1.2 mg of the title compound (160FE58E7). MS (ESI) 361 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/85.

Example 23 – 8-Chloro-4-methyl-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE74C)




[0165] 5-Chloro-2-fluoronitrobenzene (1.06 g, 6.0 mmol) and 2-amino-3-methylbenzoic acid (302 mg, 2.0 mmol) were reacted according to GP1 to give 4.8 mg of the title compound (160FE74C). MS (ESI) 327 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 97/90.

Example 24 – 1,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (203FE03)



[0166] 5-Chloro-2-fluoronitrobenzene (1.06 g, 6.0 mmol) and 2-amino-6-chlorobenzoic acid (343 mg, 2.0 mmol) were reacted according to GP1 to give 3.1 mg of the title compound (203FE03). MS (ESI) 347 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/99.

Example 25 – 8-Bromo-5-methyl-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (166JO32)



[0167] 5-Bromo-2-fluoronitrobenzene (580 mg, 2.6 mmol) and *N*-methylantranilic acid (200 mg, 1.3 mmol) were reacted according to GP1 to give 1.6 mg of the title compound (166JO32). MS (ESI) 371 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 90/74.

Example 26 – General procedure 2 (GP2)


[0168] A mixture of an aminobenzoic acid (1 eq.), a 2-fluoronitrobenzene (3 eq.) or a 2-chloronitrobenzene (3 eq.), and  $Cs_2CO_3$  (3 eq.) in DMF was heated to 140°C for 1 hour, and then allowed to obtain room temperature. The mixture was diluted with water and washed with EtOAc (2 x).

[0169] EtOH and  $Na_2S_2O_4$  (5 eq.) was added to the aqueous phase and the resulting mixture was stirred for 1 h. Aqueous HCl (2 M) was added to the mixture and then the aqueous phase was extracted with EtOAc (3 x) and the combined organic phases were concentrated.


[0170] The residue was taken up in xylene and the resulting mixture was stirred at 130°C over night. The mixture was diluted with EtOAc, washed with saturated aqueous  $NaHCO_3$ -solution, dried ( $Na_2SO_4$ ), and concentrated.

[0171] The residue was taken up in dioxane and added to a mixture of  $TiCl_4$  (1.1 eq., 1 M in toluene) and piperazine (5 eq.) in dioxane at 50°C. The resulting mixture was stirred at 100°C over night, and then allowed to obtain room temperature. Aqueous HCl (2 M) was added to the mixture until solution became acidic and then the aqueous phase was extracted with EtOAc (2 x). Aqueous NaOH (2 M) was added to the aqueous phase until a basic solution was obtained and the resulting suspension was extracted with EtOAc (3 x). The combined organic phases were concentrated and purified by HPLC.




Example 27 – 7,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (166JO28)


[0172] 1,2-Dichloro-4-fluoro-5-nitrobenzene (1.26 g, 6.0 mmol) and 2-aminobenzoic acid (274 mg, 2 mmol) were reacted according to GP2 to give 16 mg of the title compound (166JO28). MS (ESI) 347 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/96.

Example 28 – 11-(Piperazin-1-yl)-8-trifluoromethyl-5H-dibenzo[*b,e*][1,4]diazepine (166JO23)

[0173] 4-Fluoro-3-nitrobenzotrifluoride (1.25 g, 6 mmol) and 2-aminobenzoic acid (274 mg, 2 mmol) were reacted according to GP2 to give 12 mg of the title compound (166JO23). MS (ESI) 347 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 81/98.


Example 29 – 11-(Piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE19A)

[0174] 2-Fluoro-nitrobenzene (847 mg, 6.0 mmol) and 2-aminobenzoic acid (274 mg, 2 mmol) were reacted according to GP2 to give 16 mg of the title compound (160FE19A). MS (ESI) 279 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 30 – 8-Fluoro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE19C)


[0175] 2,5-Difluoronitrobenzene (955 mg, 6.0 mmol) and 2-aminobenzoic acid (274 mg, 2 mmol) were reacted according to GP2 to give 8.9 mg of the title compound (160FE19C). MS (ESI) 297 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/97.

Example 31 – 11-(Piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine-8-carbonitrile (160FE19D)




[0176] 4-Chloro-3-nitrobenzonitrile (1.10 g, 6.0 mmol) and 2-aminobenzoic acid (274 mg, 2 mmol) were reacted according to GP2 to give 4.7 mg of the title compound (160FE19D). MS (ESI) 304 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/86.

Example 32 – 8-Bromo-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE19E)



[0177] 5-Bromo-2-fluoronitrobenzene (1.32 g, 6.0 mmol) and 2-aminobenzoic acid (274 mg, 2 mmol) were reacted according to GP2 to give 15 mg of the title compound (160FE19E). MS (ESI) 357 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 33 – 8-Methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE19F)




[0178] 4-Chloro-3-nitrotoluene (1.03 g, 6.0 mmol) and 2-aminobenzoic acid (274 mg, 2 mmol) were reacted according to GP2 to give 1.6 mg of the title compound (160FE19F). MS (ESI) 293 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 70/70.

Example 34 – General procedure 3 (GP3)

[0179] 1-chloroethyl chloroformate (17 mg, 0.12 mmol) at 10°C was added to a *N*-methyl piperazine derivative (0.1 mmol) dissolved in THF (2 ml). The resulting mixture was then heated at reflux for 18 h. The temperature was decreased and the THF removed at


reduced pressure. Methanol was then added to the remaining oil and the mixture was shaken at 65°C for 2 h. The methanol was removed at reduced pressure and the remaining crude product was purified by HPLC.

Example 35 – 3-Fluoro-6-(4-methyl-piperazin-1-yl)-11*H*-dibenzo[*b,e*]azepine (160FE02)




[0180] 3-Fluoro-6-(4-methyl-piperazin-1-yl)-11*H*-dibenzo[*b,e*]azepine (31 mg, 0.1 mmol) was reacted according to GP3 to give 8 mg of the title compound isolated as oxalate salt (160FE02). MS (ESI) 296 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/100.

Example 36 – 2-(Trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE13A)




[0181] 2-(Trifluoromethanesulfonyloxy)-11-(4-methyl-piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine (39 mg, 0.1 mmol) was reacted according to GP3 to give 3.0 mg of the title compound (160FE13A). MS (ESI) 427 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 95/98.

Example 37 – 2-(Trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]oxazepine (160FE13B)




[0182] 2-(Trifluoromethanesulfonyloxy)-11-(4-methyl-piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]oxazepine (39 mg, 0.1 mmol) was reacted according to GP3 to give 11 mg of the title compound (160FE13B). MS (ESI) 428 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 98/100.

Example 38 – 8-Chloro-2-(trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE13C)



[0183] 8-Chloro-2-(trifluoromethanesulfonyloxy)-11-(4-methyl-piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (42 mg, 0.1 mmol) was reacted according to GP3 to give 3.2 mg of the title compound (160FE13C). MS (ESI) 461 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.


Example 39 – 8-(Trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE13D)




[0184] 8-(Trifluoromethanesulfonyloxy)-11-(4-methyl-piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (39 mg, 0.1 mmol) was reacted according to GP3 to give 2.2 mg of the title compound (160FE13D). MS (ESI) 427 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 40 – General procedure 4 (GP4)


[0185] A mixture of appropriate lactam (0.1 mmol) in dioxane was added to a mixture of  $TiCl_4$  (1.1 eq., 1 M in toluene) and the amine (0.5 mmol) in dioxane at 50°C or to a mixture of  $TiCl_4$  (2.2 eq., 1 M in toluene) and the amine (1.0 mmol) in dioxane at 50°C. The resulting mixture was stirred at 100°C over night, and then allowed to obtain room temperature. Aqueous HCl (3 mL, 2 M) was added to the aqueous mixture and then the aqueous phase was extracted with EtOAc (2 x 4 mL). Aqueous NaOH (6 mL, 2 M) was added to the aqueous phase and the resulting suspension was extracted with EtOAc (3 x 3mL). The combined organic phases were concentrated and purified by HPLC.

Example 41 – 11-(Piperazin-1-yl)-dibenzo[*b,f*][1,4]thiazepin (160FE17A)

[0186] 10*H*-Dibenzo[*b,f*][1,4]thiazepin-11-one (23 mg, 0.1 mmol) and piperazine (43 mg, 0.5 mmol) were reacted according to GP4 to give 3.1 mg of the title compound (160FE17A). MS (ESI) 296 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 97/90.


Example 42 - 11-(Piperazin-1-yl)-2,3-dihydro-1,4-benzodioxino[6,7-*b*][1,4]benzothiazepin (160FE17B)

[0187] 2,3-Dihydro-1,4-benzodioxino[6,7-*b*][1,4]benzothiazepin-11(12*H*)-one (29 mg, 0.1 mmol) and piperazine (43 mg, 0.5 mmol) were reacted according to GP4 to give 1.9 mg of the title compound (160FE17B). MS (ESI) 354 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/95.

Example 43 – 8-Chloro-11-[1,4]diazepam-1-yl-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE16A)


[0188] 8-Chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (25 mg, 0.1 mmol) and homopiperazine (50 mg, 0.5 mmol) were reacted according to GP4 to give 12 mg of the title compound (160FE16A). MS (ESI) 327 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/93.

Example 44 – *N*'-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)-*N,N*-dimethyl-ethane-1,2-diamine (160FE16D)




**[0189]** 8-Chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (25 mg, 0.1 mmol) and *N,N*-dimethylethylenediamine (44 mg, 0.5 mmol) were reacted according to GP4 to give 20 mg of the title compound (160FE16D). MS (ESI) 315 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 45 – *N*'-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)-*N,N*-diethyl-ethane-1,2-diamine (160FE16E)



**[0190]** 8-Chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (25 mg, 0.1 mmol) and *N,N*-diethylethylenediamine (58 mg, 0.5 mmol) were reacted according to GP4 to give 3.9 mg of the title compound (160FE16E). MS (ESI) 343 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/94.


Example 46 – 8-Chloro-11-(4-methyl-[1,4]diazepam-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE16F)



**[0191]** 8-Chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (25 mg, 0.1 mmol) and 1-methylhomopiperazine (57 mg, 0.5 mmol) were reacted according to GP4 to


give 5.7 mg of the title compound (160FE16F). MS (ESI) 341 ( $\text{MH}^+$ ). Purity for  $\text{MH}^+$  (UV/MS) 100/100.

Example 47 – 8-Chloro-2-methoxy-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE20A)




[0192] 8-Chloro-2-methoxy-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (28 mg, 0.1 mmol) and piperazine (86 mg, 1.0 mmol) were reacted according to GP4 to give 19 mg of the title compound (160FE20A). MS (ESI) 342 ( $\text{MH}^+$ ). Purity for  $\text{MH}^+$  (UV/MS) 99/100.

Example 48 – *N'*-(5H-Dibenzo[*b,e*][1,4]diazepine-11-yl)-*N,N*-dimethyl-ethane-1,2-diamine (160FE20B)




[0193] 5,10-Dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (160FE15A) (21 mg, 0.1 mmol) and *N,N*-dimethylethylenediamine (88 mg, 1.0 mmol) were reacted according to GP4 to give 7.6 mg of the title compound (160FE20B). MS (ESI) 281 ( $\text{MH}^+$ ). Purity for  $\text{MH}^+$  (UV/MS) 100/100.

Example 49 – 11-[1,4]Diazepam-1-yl-5H-dibenzo[*b,e*][1,4]diazepine (160FE20C)




[0194] 5,10-Dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (160FE15A) (21 mg, 0.1 mmol) and homopiperazine (100 mg, 1.0 mmol) were reacted according to GP4 to give 12 mg of the title compound (160FE20C). MS (ESI) 293 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 95/95.

Example 50 – *N'*-(8-Fluoro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)-*N,N*-dimethyl-ethane-1,2-diamine (160FE20D)



[0195] 8-Fluoro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (160FE15C) (23 mg, 0.1 mmol) and *N,N*-dimethylethylenediamine (88 mg, 1.0 mmol) were reacted according to GP4 to give 11 mg of the title compound (160FE20D). MS (ESI) 299 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.


Example 51 – 8-Fluoro-11-[1,4]diazepam-1-yl-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE16A)



[0196] 8-Fluoro-5,10-Dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (160FE15C) (23 mg, 0.1 mmol) and homopiperazine (100 mg, 1.0 mmol) were reacted according to GP4 to give 19 mg of the title compound (160FE20E). MS (ESI) 311 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.




Example 52 – N'-(8-Chloro-5H-dibenzo[*b,e*][1,4]diazepine-11-yl)-N-methyl-ethane-1,2-diamine (160FE22)




[0197] 8-Chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (25 mg, 0.1 mmol) and *N*-methylethylenediamine (74 mg, 1.0 mmol) were reacted according to GP4 to give 7.6 mg of the title compound (160FE22). MS (ESI) 301 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 92/83.

Example 53 – 8-Chloro-11-(trans-2,5-dimethyl-piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE33A)




[0198] 8-Chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (25 mg, 0.1 mmol) and trans-2,5-dimethylpiperazine (114 mg, 1.0 mmol) were reacted according to GP4 to give 1.9 mg of the title compound (160FE33A). MS (ESI) 341 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 100/82.

Example 54 – 8-Chloro-11-(3,5-dimethyl-piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE33B)




[0199] 8-Chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (25 mg, 0.1 mmol) and 2,6-dimethylpiperazine (114 mg, 1.0 mmol) were reacted according to GP4 to give 18 mg of the title compound (160FE33B). MS (ESI) 341 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 55 – 8-Chloro-11-(3-methyl-piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE38)




[0200] 8-Chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (25 mg, 0.1 mmol) and 2-methylpiperazine (100 mg, 1.0 mmol) were reacted according to GP4 to give 30 mg of the title compound (160FE38). MS (ESI) 327 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/89.

Example 56 – 8-Chloro-11-(3-phenyl-piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE45)




[0201] 8-Chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (25 mg, 0.1 mmol) and 2-phenylpiperazine (162 mg, 1.0 mmol) were reacted according to GP4 to give 27 mg of the title compound (160FE45). MS (ESI) 389 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/89.

Example 57 – 8-Chloro-5-methyl-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (189JO25A)



[0202] NaH (12 mg, 0.29 mmol, 60 % in mineral oil) was added to a mixture of 8,5-dichloro-5H-dibenzo[*b,e*][1,4]diazepine (160FE64) (50 mg, 0.19 mmol) in toluene (1.5 mL) and DMF (0.5 mL). MeI (24  $\mu$ L, 0.38 mmol) was then added. The resulting mixture was stirred for 1 h then quenched by addition of saturated aqueous NaHCO<sub>3</sub>-solution (2 mL). The mixture was extracted with diethyl ether, and the combined organic phases were dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. The residue was taken up in toluene (2.0 mL), piperazine (98 mg, 1.1 mmol) was added, and the resulting mixture was stirred at 100°C for 1 h. Aqueous HCl (1 mL, 2M) and EtOAc (2 mL) was then added to the mixture. The phases were separated and the aqueous phase was extracted with EtOAc (2 mL) and then aqueous NaOH (2 mL, 2 M) was added. The basic aqueous phase was extracted with EtOAc (3 x 2 mL) and the combined organic phases were dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. The residue was dissolved in DMF and purified on HPLC to give 34 mg of the title compound (189JO25A). MS (ESI) 327 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 100/100.


Example 58 – 8-Chloro-5-benzyl-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE46-PIPBN)



[0203] 8,5-Dichloro-5H-dibenzo[*b,e*][1,4]diazepine (160FE46) (51 mg, 0.20 mmol) and benzyl bromide (68 mg, 0.4 mmol) were reacted as described for Example 57 to


give 8.4 mg of the title compound (160FE46-PIPBN). MS (ESI) 403 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 59 – 8-Iodo-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (166JO38)



**[0204]** A mixture of 8-bromo-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (166JO31) (60 mg, 0.21 mmol), NaI, (62 mg, 0.42 mmol), *N,N*-dimethylethylenediamine (2.2  $\mu$ L, 0.021 mmol) and CuI (2 mg, 0.01 mmol) in dioxane (1 mL) was heated in a capped tube for 3 days. The reaction mixture was allowed to obtain room temperature and then the mixture was applied onto a SCX-2 ion exchange column and the product was eluted with  $CH_2Cl_2$  to give 49 mg of intermediate 8-iodolactam. The intermediate 8-iodolactam (20 mg, 0.060 mmol) in dioxane (1 mL) was added to a mixture of  $TiCl_4$  (0.13 mL, 0.13 mmol, 1 M in toluene) and piperazine (0.051 g, 0.60 mmol) in dioxane at 50°C. The resulting mixture was stirred at 100°C over night then allowed to obtain room temperature. Aqueous HCl (3 mL, 2 M) was added to the mixture and then the aqueous phase was extracted with EtOAc (2 x 4 mL). Aqueous NaOH (6 mL, 2 M) was added to the aqueous phase and the resulting suspension was extracted with EtOAc (3 x 3mL). The combined organic phases were concentrated and purified by HPLC to give 4.1 mg of the title compound (166JO38). MS (ESI) 405 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.


Example 60 – 2-Iodo-8-chloro-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (166JO54)



**[0205]** 2-Bromo-8-chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (intermediate from GP1) (30 mg, 0.09 mmol) was reacted as described for Example 59 to


give 7.0 mg of the title compound (166JO54). MS (ESI) 439 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 61 – 8-Phenyl-11-(piperazin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (189JO53)




[0206] Tetrakis(triphenylphosphine)palladium(0) (catalytic amount) was added to a mixture of 8-bromo-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (166JO31) (30 mg, 0.12 mmol), benzene boronic acid (18 mg, 0.15 mmol) and  $K_2CO_3$  (34 mg, 0.24 mmol) in deoxygenised toluene/EtOH/ $H_2O$  (1.5 mL) and the resulting mixture was stirred at 80°C over night. The mixture was diluted with EtOAc, washed with saturated aqueous  $NaHCO_3$ -solution, dried ( $Na_2SO_4$ ) and concentrated to give crude 8-phenyl lactam. The intermediate 8-phenyl lactam in dioxane (1 mL) was added to a mixture of  $TiCl_4$  (0.24 mL, 0.24 mmol, 1 M in toluene) and piperazine (0.103 g, 1.2 mmol) in dioxane at 50°C. The resulting mixture was stirred at 100°C over night, and then allowed to obtain room temperature. Aqueous HCl (3 mL, 2 M) was added to the mixture and then the aqueous phase was extracted with EtOAc (2 x 4 mL). Aqueous NaOH (6 mL, 2 M) was added to the aqueous phase and the resulting suspension was extracted with EtOAc (3 x 3mL). The combined organic phases were applied onto a SCX-2 ion exchange column. The column was washed with MeOH, and then the product was eluted with  $NH_3$  (7 N in MeOH), concentrated, and purified by HPLC to give 16 mg of the title compound (189JO53). MS (ESI) 355 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 62 – 8-Chloro-11-(piperidin-1-yl)-5H-dibenzo[*b,e*][1,4]diazepine (166JO69A)




[0207] Piperidine (37 mg, 0.44 mmol) was added to crude 8-chloro-11-methylsulfanyl-5*H*-dibenzo[*b,e*][1,4]diazepine(166JO50) (90 mg, purity 50%, 0.218 mmol) in pyridine (2 mL) and the resulting mixture was heated in a capped tube at 160°C for 10 h. The mixture was concentrated and flash chromatographed (SiO<sub>2</sub>, heptane:EtOAc 8:1-6:1) to give 12 mg of the title compound (166JO69A). MS (ESI) 312 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 100/100.

Example 63 – 8-Chloro-11-(morpholin-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine (166JO69B)



[0208] Crude 8-chloro-11-methylsulfanyl-5*H*-dibenzo[*b,e*][1,4]diazepine(166JO50) (90 mg, purity 50%, 0.218 mmol) and morpholine (38 mg, 0.44 mmol) were reacted as described for Example 62 to give 11 mg of the title compound (166JO69B). MS (ESI) 314 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 100/98.

Example 64 – 5-Allyl-8-chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine (166JO68)




[0209] K<sup>t</sup>OBu (343 mg, 3.1 mmol) was added to a mixture of 8-chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (500 mg, 2.0 mmol) in dioxane (10 mL) and the resulting mixture was stirred at 60°C for 1 h, then cooled to room temperature. *p*-Methoxybenzyl chloride (0.42 mL, 3.1 mmol) was added and the resulting mixture was stirred at 40°C for 2h. The reaction was quenched by addition of MeOH (2 mL). The mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed with saturated aqueous NaHCO<sub>3</sub>-solution, dried

(Na<sub>2</sub>SO<sub>4</sub>), concentrated and flash chromatographed (SiO<sub>2</sub>, heptane:EtOAc, 4:1-3:1) which gave intermediate *p*-methoxybenzylprotected lactam (732 mg), 85 % pure, which was used in the next step without further purification.

[0210] To a mixture of *p*-methoxybenzylprotected lactam (100 mg, 0.27 mmol) in DMF (2 mL) was added NaH (16 mg, 0.41 mmol, 60 % in mineral oil) and the resulting mixture was heated to 60°C then allowed to obtain room temperature. Allyl bromide (36 µL, 0.41 mmol) was added and the resulting mixture was stirred at room temperature for 3 h then diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed with saturated aqueous NaHCO<sub>3</sub>-solution, dried (Na<sub>2</sub>SO<sub>4</sub>), concentrated, flash chromatographed (SiO<sub>2</sub>, heptane:EtOAc 8:1-4:1), and concentrated. The residue was taken up in trifluoroacetic acid (4 mL) and the resulting mixture was stirred at room temperature over night, then at 45°C for 2h. The mixture was concentrated, chromatographed (SiO<sub>2</sub>, heptane:EtOAc 8:1-4:1), and concentrated. The residue was taken up in toluene (2 mL) and *N,N*-dimethylaniline (48 µL, 0.38 mmol) and POCl<sub>3</sub> (35 µL, 0.38 mmol) were added. The resulting mixture was stirred at 100°C for 2 h then concentrated. The residue was taken up in dioxane, piperazine (65 mg, 0.76 mmol) was added and the resulting mixture was stirred at 100°C for 3h. To the mixture was added aqueous HCl (3 mL, 2 M) and then the aqueous phase was extracted with EtOAc (2 x 4 mL). To the aqueous phase was added aqueous NaOH (6 mL, 2 M) and the resulting suspension was extracted with EtOAc (3 x 3 mL). The combined organic phases were concentrated and purified by HPLC to give 17 mg of the title compound (166JO68). MS (ESI) 353 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 99/88.

Example 65 – 6-Chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine (189JO68)




[0211] A mixture of a methyl 2-aminobenzoate (454 mg, 3.0 mmol), 3-chloro-2-fluoronitrobenzene (352 mg, 2 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (0.78 g, 2.4 mol) in DMF (4 mL) was stirred at 140°C for 2 h.

[0212] The mixture was diluted with EtOAc (10 mL) and washed with 2 M aqueous NaOH-solution (2 x 5 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), concentrated and flash chromatographed (SiO<sub>2</sub>, toluene:heptane:EtOAc-system) and concentrated. The residue was taken up in THF (10 mL), 1 M aqueous LiOH (5 mL) was added and the resulting mixture was stirred at 80°C for 1 h, and then allowed to obtain room temperature. 2 M aqueous HCl was added until pH 2. The aqueous phase was extracted with EtOAc (3 x). The combined organic phases were dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. The residue was taken up in EtOH and a mixture of K<sub>2</sub>CO<sub>3</sub> (1.38 g, 10 mmol) and Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> (1.74 g, 10 mmol) in water was added and the resulting mixture was stirred for 1 h. The mixture was diluted with water and washed with 1 M aqueous NaOH-solution (2 x 5 mL) and then dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated.

[0213] The residue was taken up in CH<sub>2</sub>Cl<sub>2</sub> and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (307 mg, 1.6 mmol) was added. The resulting mixture was stirred at room temperature for 1 h. The mixture was diluted with EtOAc, washed with saturated aqueous NaHCO<sub>3</sub>-solution, dried (Na<sub>2</sub>SO<sub>4</sub>), concentrated, and flash chromatographed (SiO<sub>2</sub>, heptane:EtOAc, 2:1) to give 21 mg of the intermediate lactam.

[0214] The intermediate lactam was taken up in dioxane and added to a mixture of TiCl<sub>4</sub> (0.19 mL, 0.19 mmol, 1 M in toluene) and piperazine (73 mg, 0.85 mmol) in dioxane at 50°C. The resulting mixture was stirred at 100°C over night, and then allowed to obtain room temperature. To the mixture was added aqueous HCl (1 mL, 2 M) and then the aqueous phase was extracted with EtOAc (2 x 1 mL). To the aqueous phase was added aqueous NaOH (2 mL, 2 M) and the resulting suspension was extracted with EtOAc (3 x 1 mL). The combined organic phases were concentrated and purified by HPLC to give 9.8 mg of the title compound (189JO68) MS (ESI) 313 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 100/98.


Example 66 – 8-Chloro-5-piperazin-1-yl-11H-benzo[b]pyrido[2,3-e][1,4]diazepine (166JO63)





[0215] To a mixture of 5-chloro-2-nitroaniline (345 mg, 2 mmol) and pyridine (162  $\mu$ L, 2 mmol) in dioxane was added 2-chloronicotinyl chloride (352 mg, 2 mmol) and the resulting mixture was stirred at room temperature for 2h. The mixture was diluted with  $\text{CH}_2\text{Cl}_2$ , washed with saturated aqueous  $\text{NaHCO}_3$ -solution, dried ( $\text{Na}_2\text{SO}_4$ ), concentrated and crystallised from MeOH to give 271 mg of intermediate diarylamine. To a mixture of intermediate diarylamine (100 mg, 0.32 mmol) in EtOH (0.5 mL) was added a mixture of  $\text{K}_2\text{CO}_3$  (220 mg, 1.6 mmol) and  $\text{Na}_2\text{S}_2\text{O}_4$  (278 mg, 1.6 mmol) in water (0.5 mL) and the resulting mixture was stirred for 1 h at room temperature. The mixture was concentrated and the residue taken up in EtOAc/ $\text{H}_2\text{O}$  and separated. The organic phase was dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated. The residue was taken up in xylene and heated to  $130^\circ\text{C}$  over night, then diluted with EtOAc, washed with saturated aqueous  $\text{NaHCO}_3$ -solution, dried ( $\text{Na}_2\text{SO}_4$ ), concentrated and flash chromatographed ( $\text{SiO}_2$ , heptane:EtOAc) to give intermediate lactam. The intermediate lactam was taken up in dioxane and added to a mixture of  $\text{TiCl}_4$  (187  $\mu$ L, 0.187 mmol, 1 M in toluene) and piperazine (73 mg, 0.85 mmol) in dioxane at  $50^\circ\text{C}$ . The resulting mixture was stirred at  $100^\circ\text{C}$  over night, and then allowed to obtain room temperature. To the mixture was added aqueous HCl (1 mL, 2 M) and then the aqueous phase was extracted with EtOAc (2 x 2 mL). To the aqueous phase was added aqueous NaOH (2 mL, 2 M) and the resulting suspension was extracted with EtOAc (3 x 1 mL). The combined organic phases were concentrated and purified by HPLC to give 20 mg of the title compound (166JO63). MS (ESI) 314 ( $\text{MH}^+$ ). Purity for  $\text{MH}^+$  (UV/MS) 100/99.

Example 67 – 2-Chloro-10-piperazin-1-yl-5H-dibenzo[*b,f*]azepin (189JO39)




[0216] To a mixture under Ar of 2-chloro-5-(4-methoxybenzyl)-5,11-dihydrodibenzo[*b,f*]azepin-11-one (189JO27) (150 mg, 0.41 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) at  $-75^\circ\text{C}$  was added  $\text{TiCl}_4$  (0.60 mL, 0.60 mmol, 1 M in toluene) and the resulting mixture was stirred for 1 h. The mixture was diluted with saturated aqueous  $\text{NH}_4\text{Cl}$ -solution and  $\text{CH}_2\text{Cl}_2$


and the mixture was allowed to obtain room temperature and the phases were separated. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> (1 x 10 mL) and the combined organic phases were dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to give crude protected product (90 mg, 90%), that was used in the next step without further purification.

[0217] To a solution of TiCl<sub>4</sub> (0.18 mL, 0.18 mmol, 1 M in toluene) and piperazine (283 mg, 3.3 mmol) in dioxane (4 mL) at 50° was added crude protected product (80 mg, 0.33 mmol) and the resulting suspension was stirred at 100°C for 1.5 h. The mixture was allowed to obtain room temperature, then it was diluted with EtOAc, washed with saturated aqueous NaHCO<sub>3</sub>-solution, dried (Na<sub>2</sub>SO<sub>4</sub>), concentrated and flash chromatographed (Al<sub>2</sub>O<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>:MeOH, 1:0-25:1) to give 64 mg (63%) of the title compound (189JO39). MS (ESI) 312 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 97/95.


Example 68 – 8-Chloro-11-(piperazin-1-yl)-dibenzo[b,f][1,4]thiazepine (189JO16)



[0218] To a mixture of 8-chloro-10*H*-dibenzo[b,f][1,4]thiazepin-11-one (189JO13) (38 mg, 0.15 mmol) and *N,N*-dimethylaniline (46 μL, 0.36 mmol) in toluene was added POCl<sub>3</sub> (27 μL, 0.29 mmol) and the resulting mixture was stirred for 2 h at 100°C, and then concentrated. Toluene (2 mL) and piperazine (62 mg, 0.73 mmol) were added, and the resulting mixture was stirred at 100°C for 3 h, and then allowed to obtain room temperature. To the mixture was added aqueous HCl (1 mL, 2 M) and then the aqueous phase was extracted with EtOAc (2 x 2 mL). To the aqueous phase was added aqueous NaOH (3 mL, 2 M) and the resulting mixture was extracted with EtOAc (3 x 3 mL). The combined organic phases were concentrated and purified by HPLC to give 6.6 mg of the title compound (189JO16). MS (ESI) 330 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 99/98.

Example 68 – 8-Chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO31)


[0219] A mixture of 8-chloro-10*H*-dibenzo[*b,f*][1,4]oxazepin-11-one (189JO29C) (17 mg, 0.069 mmol) and 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (16 mg, 0.040 mmol) in toluene (2 mL) was heated in capped tube using microwave assisted heating (130°C, 20 minutes). The reaction mixture was cooled to room temperature and MeI (18  $\mu$ L, 0.29 mmol) was added and the resulting mixture was heated in capped tube using microwave assisted heating (120°C, 20 minutes). The mixture was concentrated and the residue was taken up in pyridine (2 mL) and piperazine (25 mg, 0.29 mmol) was added. The resulting mixture was heated in a capped tube at 130°C over night then using microwave assisted heating (160°C, 30 minutes). The mixture was concentrated, diluted with EtOAc and washed with water. The organic phase was applied onto a SCX-2 ion exchange column. The column was washed with MeOH, and then the product was eluted with NH<sub>3</sub> (7 N in MeOH) to give 9.0 mg (57%) of the title compound (189JO31). MS (ESI) 314 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 92/100.

Example 69 – 8-Chloro-11-(4-methyl-piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO47)

[0220] A mixture of 8-chloro-10*H*-dibenzo[*b,f*][1,4]oxazepin-11-one (189JO29C) (30 mg, 0.069 mmol) and 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (29 mg, 0.040 mmol) in toluene (2 mL) was heated in a capped tube using microwave assisted heating (130°C, 20 minutes). The reaction mixture was

cooled to room temperature and MeI (38  $\mu$ L, 0.29 mmol) was added and the resulting mixture was heated in capped tube using microwave assisted heating (120°C, 20 minutes). The mixture was concentrated and the residue was taken up in pyridine (2 mL) and piperazine (24 mg, 0.29 mmol) was added. The resulting mixture was heated in a capped tube at 130°C over night then heated using microwave assisted heating (160°C, 30 minutes). The mixture was concentrated, diluted with EtOAc and washed with water. The organic phase was dried ( $\text{Na}_2\text{SO}_4$ ), concentrated and flash chromatographed ( $\text{SiO}_2$ , toluene:EtOAc:MeOH, 4:2:0-2:2:1) to give 8.9 mg of the title compound (189JO47). MS (ESI) 328 ( $\text{MH}^+$ ). Purity for  $\text{MH}^+$  (UV/MS) 98/93.

Example 70 – 3-Chloro-6-piperazin-1-yl-11H-dibenzo[b,e]azepine (189JO60)



[0221] 3-Chloro-5,11-dihydro-dibenzo[b,e]azepin-6-one (189JO59) (25 mg, 0.1 mmol) and piperazine were reacted according to GP4 to give 2.2 mg of the title compound (189JO60). MS (ESI) 312 ( $\text{MH}^+$ ). Purity for  $\text{MH}^+$  (UV/MS) 100/100.

Example 71 – General procedure 5 (GP5)


[0222] A mixture of a methyl aminobenzoic ester (2.0 mmol), a 2-fluoronitrobenzene (1.0 mmol) and  $\text{Cs}_2\text{CO}_3$  (0.65 g, 2.0 mmol) in DMF (4mL) was stirred at 40°C for 2h. The mixture was diluted with EtOAc (10 mL) and washed with 2 M aqueous NaOH-solution (2 x 5 mL).

[0223] EtOH,  $\text{H}_2\text{O}$ ,  $\text{K}_2\text{CO}_3$  (0.69 g, 5 mmol) and  $\text{Na}_2\text{S}_2\text{O}_4$  (0.87 g, 5 mmol) was added to the EtOAc-phase and the resulting mixture was stirred vigorously for 1 h. The aqueous phase was removed and the organic phase was washed with 1 M aqueous NaOH-solution (2 x 5 mL) and then concentrated.

[0224] The residue was taken up in DMF (1 mL), toluene (4 mL) and NaH (60 mg, 1.5 mmol, 60% in mineral oil) was added and the resulting mixture was stirred at 80°C over night, then quenched by addition of saturated aqueous  $\text{NH}_4\text{Cl}$ -solution. The resulting mixture was diluted with EtOAc, washed with 2 M aqueous NaOH-solution (2 x 5 mL),


dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated. The residue was taken up in dioxane and added to a mixture of  $\text{TiCl}_4$  (1.1 mL, 1.1 mmol, 1 M in toluene) and piperazine (0.41 g, 5 mmol) in dioxane at  $50^\circ\text{C}$ . The resulting mixture was stirred at  $100^\circ\text{C}$  over night, and then allowed to obtain room temperature. To the mixture was added aqueous  $\text{HCl}$  (3 mL, 2 M) and then the aqueous phase was extracted with  $\text{EtOAc}$  (2 x 4 mL). To the aqueous phase was added aqueous  $\text{NaOH}$  (6 mL, 2 M) and the resulting suspension was extracted with  $\text{EtOAc}$  (3 x 3 mL). The combined organic phases were concentrated, dried ( $\text{Na}_2\text{SO}_4$ ) and purified by HPLC.

Example 72 – 8-Bromo-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO48A)




[0225] 5-Bromo-2-fluoronitrobenzene (220 mg, 1 mmol) and methyl 2-hydroxybenzoate (304 mg, 2 mmol) were reacted according to GP5 to give 36 mg of the title compound (189JO48A). MS (ESI) 358 ( $\text{MH}^+$ ). Purity for  $\text{MH}^+$  (UV/MS) 96/82.

Example 73 – 11-(Piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO48B)




[0226] 2-fluoronitrobenzene (141 mg, 1 mmol) and methyl 2-hydroxybenzoate (304 mg, 2 mmol) were reacted according to GP5 to give 5.2 mg of the title compound (189JO48B). MS (ESI) 280 ( $\text{MH}^+$ ). Purity for  $\text{MH}^+$  (UV/MS) 99/99.

Example 74 – 7-Chloro-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO50A)




[0227] 4-Chloro-2-fluoronitrobenzene (175 mg, 1 mmol) and methyl 2-hydroxybenzoate (304 mg, 2 mmol) were reacted according to GP5 to give 17 mg of the title compound (189JO50A). MS (ESI) 314 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 75 – 8-Chloro-3-methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO50B)




[0228] 5-Chloro-2-fluoronitrobenzene (175 mg, 1 mmol) and methyl 2-hydroxy-4-methoxybenzoate (364 mg, 2 mmol) were reacted according to GP5 to give 6.8 mg of the title compound (189JO50B). MS (ESI) 344 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 94/86.


Example 76 – 8-Bromo-3-methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO50D)




[0229] 5-Bromo-2-fluoronitrobenzene (220 mg, 1 mmol) and methyl 2-hydroxy-4-methoxybenzoate (364 mg, 2 mmol) were reacted according to GP5 to give 14 mg of the title compound (189JO50D). MS (ESI) 388 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 77 – 3-Methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO50E)

[0230] 2-Fluoronitrobenzene (141 mg, 1 mmol) and methyl 2-hydroxy-4-methoxybenzoate (364 mg, 2 mmol) were reacted according to GP5 to give 33 mg of the title compound (189JO50E). MS (ESI) 310 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.


Example 78 – 7-Chloro-3-methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO50F)

[0231] 4-Chloro-2-fluoronitrobenzene (175 mg, 1 mmol) and methyl 2-hydroxy-4-methoxybenzoate (364 mg, 2 mmol) were reacted according to GP5 to give 6.7 mg of the title compound (189JO50F). MS (ESI) 344 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 98/96.

Example 79 – 8-Chloro-4-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO50H)


[0232] 5-Chloro-2-fluoronitrobenzene (175 mg, 1 mmol) and methyl 2-hydroxy-3-methylbenzoate (332 mg, 2 mmol) were reacted according to GP5 to give 34 mg of the title compound (189JO50H). MS (ESI) 328 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 80 – 8-Bromo-4-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO51A)




[0233] 5-Bromo-2-fluoronitrobenzene (220 mg, 1 mmol) and methyl 2-hydroxy-3-methylbenzoate (332 mg, 2 mmol) were reacted according to GP5 to give 20 mg of the title compound (189JO51A). MS (ESI) 372 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 81 – 4-Methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO51B)




[0234] 2-Fluoronitrobenzene (141 mg, 1 mmol) and methyl 2-hydroxy-3-methylbenzoate (332 mg, 2 mmol) were reacted according to GP5 to give 1.8 mg of the title compound (189JO51B). MS (ESI) 294 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/98.

Example 82 – 2-Bromo-8-chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO51D)




[0235] 5-Chloro-2-fluoronitrobenzene (175 mg, 1 mmol) and methyl 5-bromo-2-hydroxybenzoate (462 mg, 2 mmol) were reacted according to GP5 to give 21 mg of the title compound (189JO51D). MS (ESI) 392 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.




Example 83 – 2,8-Dibromo-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO51E)

[0236] 5-Bromo-2-fluoronitrobenzene (220 mg, 1 mmol) and methyl 5-bromo-2-hydroxybenzoate (462 mg, 2 mmol) were reacted according to GP5 to give 0.7 mg of the title compound (189JO51E). MS (ESI) 436 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 94/99.


Example 84 – 2-Bromo-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO51F)

[0237] 2-Fluoronitrobenzene (142 mg, 1 mmol) and methyl 5-bromo-2-hydroxybenzoate (462 mg, 2 mmol) were reacted according to GP5 to give 10 mg of the title compound (189JO51F). MS (ESI) 358 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 95/99.

Example 85 – 2-Bromo-7-chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO51G)


[0238] 4-Chloro-2-fluoronitrobenzene (175 mg, 1 mmol) and methyl 5-bromo-2-hydroxybenzoate (462 mg, 2 mmol) were reacted according to GP5 to give 17 mg of the title compound (189JO51G). MS (ESI) 392 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 86 – 11-(Piperazin-1-yl)-8-trifluoromethyl-dibenzo[*b,f*][1,4]oxazepine (189JO54A)




[0239] 2-Fluoro-3-nitrobenzotrifluoride (209 mg, 1 mmol) and methyl 2-hydroxybenzoate (304 mg, 2 mmol) were reacted according to GP5 to give 19 mg of the title compound (189JO54A). MS (ESI) 348 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 100/100.

Example 87 – 4-Methyl-11-(piperazin-1-yl)-8-trifluoromethyl-dibenzo[*b,f*][1,4]oxazepine (189JO54C)




[0240] 2-Fluoro-3-nitrobenzotrifluoride (209 mg, 1 mmol) and methyl 2-hydroxy-3-methylbenzoate (332 mg, 2 mmol) were reacted according to GP5 to give 15 mg of the title compound (189JO54C). MS (ESI) 362 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 100/100.

Example 88 – 8-Fluoro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO54E)




[0241] 2,5-Difluoronitrobenzene (159 mg, 1 mmol) and methyl 2-hydroxybenzoate (304 mg, 2 mmol) were reacted according to GP5 to give 14 mg of the title compound (189JO54E). MS (ESI) 298 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 100/100.

Example 89 – 8-Fluoro-3-methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO54F)




[0242] 2,5-Difluoronitrobenzene (159 mg, 1 mmol) and methyl 2-hydroxy-4-methoxybenzoate (364 mg, 2 mmol) were reacted according to GP5 to give 9.8 mg of the title compound (189JO54F). MS (ESI) 328 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 90 – 8-Fluoro-4-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO54G)




[0243] 2,5-Difluoronitrobenzene (159 mg, 1 mmol) and 2-hydroxy-3-methylbenzoate (332 mg, 2 mmol) were reacted according to GP5 to give 9.8 mg of the title compound (189JO54G). MS (ESI) 312 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.


Example 91 – 2-Bromo-8-fluoro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO54H)




[0244] 2,5-Difluoronitrobenzene (159 mg, 1 mmol) and methyl 5-bromo-2-hydroxybenzoate (462 mg, 2 mmol) were reacted according to GP5 to give 11 mg of the title compound (189JO54H). MS (ESI) 376 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 92 – 8-Methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO58A)

[0245] 4-Fluoro-3-nitrotoluene (155 mg, 1 mmol) and methyl 2-hydroxybenzoate (304 mg, 2 mmol) were reacted according to GP5 to give 24 mg of the title compound (189JO58A). MS (ESI) 294 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/98.


Example 93 – 3-Methoxy-8-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO58B)

[0246] 4-Fluoro-3-nitrotoluene (155 mg, 1 mmol) and methyl 2-hydroxy-4-methoxybenzoate (364 mg, 2 mmol) were reacted according to GP5 to give 27 mg of the title compound (189JO58B). MS (ESI) 324 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/98.

Example 94 – 4,8-Dimethyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO58C)


[0247] 4-Fluoro-3-nitrotoluene (155 mg, 1 mmol) and methyl 2-hydroxy-3-methylbenzoate (332 mg, 2 mmol) were reacted according to GP5 to give 24 mg of the title compound (189JO58C). MS (ESI) 308 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/98.

Example 95 – 3-Methoxy-11-(piperazin-1-yl)-8-trifluoromethyl-dibenzo[*b,f*][1,4]oxazepine (189JO62A)




[0248] 2-Fluoro-3-nitrobenzotrifluoride (209 mg, 1 mmol) and methyl 2-hydroxy-4-methoxybenzoate (364 mg, 2 mmol) were reacted according to GP5 to give 12 mg of the title compound (189JO62A). MS (ESI) 378 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/95.

Example 96 – 2-Bromo-11-(piperazin-1-yl)-8-trifluoromethyl-dibenzo[*b,f*][1,4]oxazepine (189JO62B)



[0249] 2-Fluoro-3-nitrobenzotrifluoride (209 mg, 1 mmol) and methyl 5-bromo-2-hydroxybenzoate (462 mg, 2 mmol) were reacted according to GP5 to give 11 mg of the title compound (189JO62B). MS (ESI) 426 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 97 – 6-Chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO69)




[0250] 3-Chloro-2-fluoronitrobenzene (352 mg, 2 mmol) and methyl 2-hydroxybenzoate (453 mg, 3 mmol) were reacted according to GP5 to give 57 mg of the title compound (189JO69). MS (ESI) 314 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 98 – General procedure 6 (GP6)


[0251] A mixture of a methyl aminobenzoic ester (1.0 mmol), a 2-fluoronitrobenzene (0.5 mmol) and  $\text{Cs}_2\text{CO}_3$  (0.33 g, 1.0 mol) in DMF (3 mL) was stirred at 40°C for 2 h. The mixture was diluted with EtOAc (10 mL) and washed with 2 M aqueous NaOH-solution (2 x 5 mL), dried ( $\text{Na}_2\text{SO}_4$ ), concentrated, flash chromatographed ( $\text{SiO}_2$ , toluene:heptane:EtOAc-system), and concentrated. The residue was taken up in THF (4 mL), 1 M aqueous LiOH (3 mL) was added and the resulting mixture was stirred at 80°C for 1 h, and then allowed to obtain room temperature. 2 M aqueous HCl was added until a pH of 2 was reached. The aqueous phase was extracted with EtOAc (3 x). The combined organic phases were dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated. The residue was taken up in EtOH and a mixture of  $\text{K}_2\text{CO}_3$  (0.35 g, 2.55 mmol) and  $\text{Na}_2\text{S}_2\text{O}_4$  (0.44 g, 2.5 mmol) in water was added and the resulting mixture was stirred for 1 h. The mixture was diluted with water and washed with 1 M aqueous NaOH-solution (2 x 5 mL) and then dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated.

[0252] The residue was taken up in  $\text{CH}_3\text{CN}$ , 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (143 mg, 0.75 mmol), 1-hydroxybenzotriazole hydrate (160 mg, 0.75 mmol), triethylamine (311  $\mu\text{L}$ , 2.25 mmol), and *N,N*-dimethylaminopyridine (catalytic amount) were added. The resulting mixture was heated in a capped tube using microwave assisted heating (140°C, 10 min). The mixture was diluted with EtOAc, washed with saturated aqueous  $\text{NaHCO}_3$ -solution, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated. The residue was taken up in dioxane and added to a mixture of  $\text{TiCl}_4$  (0.55 mL, 0.55 mmol, 1 M in toluene) and piperazine (0.22 g, 2.5 mmol) in dioxane at 50°C. The resulting mixture was stirred at 100°C over night, and then allowed to obtain room temperature. To the mixture was added aqueous HCl (3 mL, 2 M) and then the aqueous phase was extracted with EtOAc (2 x 4 mL). To the aqueous phase was added aqueous NaOH (6 mL, 2 M) and the resulting suspension was extracted with EtOAc (3 x 3 mL). The combined organic phases were concentrated and purified by HPLC.

Example 99 – 2-Bromo-8-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO63A)


[0253] 4-Fluoro-3-nitrotoluene (78 mg, 0.5 mmol) and methyl 5-bromo-2-hydroxybenzoate (231 mg, 1 mmol) were reacted according to GP6 to give 13 mg of the title compound (189JO63A). MS (ESI) 372 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 100 – 7-Chloro-4-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO63B)



[0254] 4-Chloro-2-fluoronitrobenzene (88 mg, 0.5 mmol) and methyl 2-hydroxy-3-methylbenzoate (166 mg, 1 mmol) were reacted according to GP6 to give 24 mg of the title compound (189JO63B). MS (ESI) 328 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.


Example 101 – 8-Phenyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine (189JO64)



[0255] To a mixture of 8-bromo-10*H*-dibenzo[*b,f*][1,4]oxazepin-11-one (189JO56) (30 mg, 0.12 mmol), benzene boronic acid (18 mg, 0.15 mmol) and  $K_2CO_3$  (34 mg, 0.24 mmol) in deoxygenised toluene/EtOH/ $H_2O$  (1.5 mL) was added tetrakis(triphenylphosphine)palladium(0) (catalytic amount) and the resulting mixture was heated in a capped tube in a microwave oven (140°C, 15 min). The mixture was diluted with EtOAc, washed with saturated aqueous  $NaHCO_3$ -solution, dried ( $Na_2SO_4$ ) and concentrated to give crude 8-phenyl lactam. A mixture of the intermediate 8-phenyl lactam in dioxane (1 mL) was added to a mixture of  $TiCl_4$  (0.27 mL, 0.27 mmol, 1 M in toluene) and piperazine (0.103 g, 1.2 mmol) in dioxane at 50°C. The resulting mixture was stirred at 100°C over night and then allowed to obtain room temperature. To the mixture was added

aqueous HCl (3 mL, 2 M) and then the aqueous phase was extracted with EtOAc (2 x 4 mL). To the aqueous phase was added aqueous NaOH (6 mL, 2 M) and the resulting suspension was extracted with EtOAc (3 x 3 mL). The combined organic phases were applied onto a SCX-2 ion exchange column. The column was washed with MeOH, and then the product was eluted with NH<sub>3</sub> (7 N in MeOH) concentrated and purified by HPLC to give 16 mg of the title compound (189JO64). MS (ESI) 356 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 100/99.


Example 102 – 8-Chloro-11-(piperidin-4-yl)-5H-dibenzo[*b,e*][1,4]diazepine(160FE67A)



[0256] 4-CBZ-piperidylzinc iodide (generated from 4-CBZ-piperidyl iodide (345 mg, 1.0 mmol) using zinc metal and dibromoethane) (0.8 mmol) was added at 50°C to a solution of 8,5-dichloro-5H-dibenzo[*b,e*][1,4]diazepine (160FE64) (106 mg, 0.4 mmol) and PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (18 mg, 0.04 mmol) in dry THF (2 mL). The reaction was shaken for 16 h and then quenched by the addition of aqueous saturated NH<sub>4</sub>Cl-solution. The resulting mixture was extracted twice with ether and the combined ethereal phases were washed with brine and dried (Na<sub>2</sub>SO<sub>4</sub>). Filtration followed by concentration at reduced pressure of the organic phase gave a crude product. BBr<sub>3</sub> (100 µL) added at -30°C was added to the crude product dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 mL). The reaction temperature was then slowly increased to 0°C. TLC indicated complete conversion of the starting material and Et<sub>3</sub>N, H<sub>2</sub>O and EtOAc were sequentially added to the reaction mixture. The organic phase was washed with brine and dried (Na<sub>2</sub>SO<sub>4</sub>). Filtration followed by concentration at reduced pressure gave a crude product, which was purified by HPLC to give 2.3 mg of the title compound (160FE67A). MS (ESI) 312 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 99/96.



Example 103 – 5-Benzyl-8-chloro-11-(piperidin-4-yl)-5H-dibenzo[*b,e*][1,4]diazepine (160FE67B)




[0257] 4.4 mg of the title compound (160FE67B) was isolated as a by-product in the synthesis of Example 102. MS (ESI) 402 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 85/87.

Example 104 – General procedure 7 (GP7)

[0258] A mixture of a 2-aminobenzoic acid (1 eq.), a 2-fluoronitrobenzene (2 eq. or 3 eq.) and  $K_2CO_3$  (3 eq.) in DMF was heated to 100° for 2 hour then allowed to obtain room temperature. The organic phase was extracted with 0.1 M aqueous NaOH-solution (3 x). The combined aqueous phases were acidified with 4 M aqueous HCl and extracted with EtOAc (3 x). The combined organic phases were dried ( $Na_2SO_4$ ) and concentrated. The residue was taken up in EtOH and a solution of  $K_2CO_3$  (5 eq.) and  $Na_2S_2O_4$  (5 eq.) in water was added and the resulting mixture was stirred for 1 h. The mixture was concentrated and the residue taken up in EtOAc. The mixture was acidified with aqueous HCl (2 M) and then the aqueous phase was extracted with EtOAc (3 x) and the combined organic phases were concentrated.


[0259] The residue was taken up in xylene and the resulting mixture was stirred at 130°C over night. The mixture was diluted with EtOAc, washed with saturated aqueous  $NaHCO_3$ -solution, dried ( $Na_2SO_4$ ), concentrated, and flash chromatographed ( $SiO_2$ , heptane:EtOAc system)

Example 105 – 8-Bromo-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (166JO31)




**[0260]** 5-Bromo-2-fluoronitrobenzene (1.6 g, 7.4 mmol) and 2-aminobenzoic acid (0.50 g, 3.6 mmol) were reacted according to GP7 to give 331 mg of the title compound (166JO31). MS (ESI) 289 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV) 93%.

Example 106 – 5,10-Dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (160FE15A)




**[0261]** 2-Fluoronitrobenzene (847 g, 6 mmol) and 2-aminobenzoic acid (274 mg, 2.0 mmol) were reacted according to GP7 to give 130 mg of the title compound (160FE15A).

**Example 107 – 8-Fluoro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (160FE15C)**



**[0262]** 2,4-Difluoronitrobenzene (0.96 g, 6 mmol) and 2-aminobenzoic acid (274 mg, 2.0 mmol) were reacted according to GP7 to give 100 mg of the title compound (160FE15C).


Example 108 – 8,5-Dichloro-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE64)



**[0263]** *N,N*-dimethylaniline (5.1 ml, 40 mmol) and phosphorus oxychloride (2.8 ml, 30 mmol) was added to a mixture of 8-chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (2.45g, 10 mmol) in dry toluene (20 ml). The mixture was shaken at 95°C for 2h. The temperature was then decreased and the excess *N,N*-dimethylaniline and phosphorus oxychloride were removed at reduced pressure using an oil pump. The remaining oil was dissolved in dioxane (20 ml) and aqueous Na<sub>2</sub>CO<sub>3</sub>-solution (10 ml, 2 M) was added. The two-phase mixture was shaken at 80°C for 30 min. The temperature was then decreased and ether was added to the reaction mixture. The ethereal phase was washed with saturated


aqueous NaCl-solution, dried ( $\text{Na}_2\text{SO}_4$ ) and finally concentrated at reduced pressure. The obtained oil crystallized upon standing at room temperature. Recrystallization (heptane-ether) gave 1.8 g (69 %) of the title compound (160FE64).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.61 (dd, 1 H,  $J = 1.4, 7.8$  Hz), 7.31 (dt, 1 H,  $J = 1.5, 8.0$  Hz), 7.15 (d, 1 H,  $J = 2.5$  Hz), 7.02 (m 2 H), 6.66 (dd, 1 H,  $J = 1.0, 7.8$  Hz), 6.58 (d, 1 H,  $J = 8.4$  Hz), 4.94 (bs, 1H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  157.2, 152.4, 140.3, 138.9, 134.0, 131.9, 129.7, 128.5, 128.0, 127.0, 123.5, 121.0, 119.8.

Example 109 – 8-Chloro-11-methylsulfanyl-5H-dibenzo[*b,e*][1,4]diazepine(166JO50)




**[0264]** A mixture of 8-chloro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one (500 mg, 2.05 mmol) and 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (480 mg, 1.19 mmol) in toluene (4 mL) was heated in a capped tube in a microwave oven (120°C, 30 minutes). The mixture was chromatographed ( $\text{SiO}_2$ , heptane:EtOAc, 2:1) to give 599 mg of the intermediate thiolactam. To a mixture of the intermediate thiolactam in THF (10 mL) was added MeI (633  $\mu\text{L}$ , 10.3 mmol) and the resulting mixture was heated at reflux for 4 h. The mixture was concentrated to give 610 mg of the crude title compound (166JO50) (purity 50%).

Example 110 – *N,N*-diethyl(2-bromobenzyl)amide (189JO10)



**[0265]** To a mixture of 2-bromo benzoylchloride (3.5 g, 16 mmol) in  $\text{CH}_2\text{Cl}_2$  (50 mL) at 0°C was added diethylamine (3.2 mL, 32 mmol) drop-wise and the resulting mixture was allowed to obtain room temperature. After 30 minutes, water was added, the mixture was diluted with  $\text{CH}_2\text{Cl}_2$ , washed with saturated aqueous  $\text{NaHCO}_3$ -solution and saturated aqueous  $\text{NH}_4\text{Cl}$ -solution, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated to give 3.9 g (95%) of the title compound (189JO10).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.54 (m, 1 H), 7.32 (m, 1 H), 7.22 (m, 2 H), 3.79 (m, 1 H), 3.33 (m, 1 H), 3.13 (m, 2 H), 1.26 (t, 3 H,  $J = 7.2$  Hz), 1.05 (t, 3 H,  $J = 7.0$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  168.5, 139.0, 132.8, 130.0, 127.61, 127.59, 119.3, 42.8, 39.0, 14.0, 12.6


Example 111 – 2[(4-Chloro-2-methylphenyl)-(4-methoxybenzyl)-amino]-*N,N*-diethylbenzamide (189JO26)



[0266] To a mixture of *N,N*-diethyl(2-bromobenzyl)amide (189JO10) (1.41 g, 5.50 mmol) and 4-chloro-2-methylaniline (1.01 g, 7.15 mmol) in deoxygenised toluene (14 mL) was added NaO<sup>t</sup>Bu (0.74 g, 7.7 mmol), rac-BINAP (110 mg, 0.17 mmol) and Pd(OAc)<sub>2</sub> (18 mg, 0.08 mmol) and the resulting mixture was stirred under Ar for 14 h at 80°C. The mixture was filtered through celite, concentrated and flash chromatographed (SiO<sub>2</sub>, heptane:EtOAc, 10:1-4:1) which gave unprotected intermediate ketone (1.50 g) containing about 15% impurities.


[0267] The mixture containing the intermediate was dissolved in DMF (20 mL). *p*-Methoxybenzyl chloride (0.90 mL, 6.6 mmol) was added and then NaH (0.23 g, 5.6 mmol, 60% in mineral oil) was added portions-wise. The resulting mixture was stirred at room temperature for 1 h, and then quenched by addition of saturated aqueous NaHCO<sub>3</sub>-solution. The mixture was diluted with EtOAc, washed with saturated aqueous NaHCO<sub>3</sub>-solution, dried (Na<sub>2</sub>SO<sub>4</sub>), concentrated and flash chromatographed (SiO<sub>2</sub>, toluene: EtOAc 10:1) to give 1.66 g (68%) of the title compound (189JO26). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.35 (m, 2 H), 7.20 (m, 1 H), 7.09-6.99 (m, 4 H), 6.91 (m, 2 H), 6.80 (m, 2 H), 4.84 /4.54 (Abq, 2 H, *J* = 16.2 Hz), 3.74 (s, 3H), 3.18 (m, 2H), 3.03 (m, 1 H), 2.48 (m, 1 H), 2.17 (s, 3 H), 1.01 (t, 3 H, *J* = 7.2 Hz), 0.97 (t, 3 H, *J* = 7.0 Hz), <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 169.6, 158.7, 146.53, 146.51, 137.0, 131.3, 130.9, 130.4, 129.6, 129.3, 128.7, 127.8, 127.4, 126.3, 122.8, 121.4, 114.0, 57.1, 55.3, 43.3, 39.0, 19.1, 13.9, 12.9. MS (ESI) 437 (MH<sup>+</sup>).

Example 112 – 2-Chloro-5-(4-methoxybenzyl)-5,11-dihydrodibenzo[b,f]azepin-11-one (189JO27)



[0268] To a mixture of diisopropylamine (1.09 mL, 7.8 mmol) and *N,N,N,N*-tetramethylethylenediamine (1.17 mL, 7.8 mmol) in dry THF (19 mL) at  $-20^{\circ}\text{C}$  was added *n*-BuLi (5.54 mL, 1.4 M in hexane) and the resulting mixture was stirred at  $-20^{\circ}\text{C}$  for 5 minutes. Then a mixture of 2[(4-chloro-2-methylphenyl)-(4-methoxybenzyl)-amino]-*N,N*-diethylbenzamide (189JO26) (1.36 g, 3.1 mmol) in dry THF (38 mL) was added and the resulting mixture was stirred at  $-20^{\circ}$  for 4 h. The reaction was quenched by addition of saturated aqueous  $\text{NH}_4\text{Cl}$ -solution. The mixture was diluted with EtOAc, washed with water, dried ( $\text{Na}_2\text{SO}_4$ ), concentrated, and flash chromatographed ( $\text{SiO}_2$ , toluene:heptane, 7:1-1:0) to give 665 mg (59%) of the title compound (189JO27).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  8.15 (dd, 1 H,  $J = 1.8, 8.0$  Hz), 7.43 (m, 1 H), 7.24 (m, 4 H), 7.17 (d, 1 H,  $J = 8.6$  Hz), 7.12 (dd, 1 H,  $J = 2.4, 8.6$  Hz), 7.00 (dt, 1 H,  $J = 0.8, 7.0$  Hz), 6.81 (m, 2 H), 5.09 (s, 2 H), 4.00 (s, 2 H), 3.75 (s, 3 H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  190.3, 159.1, 149.5, 146.2, 134.1, 132.4, 131.3, 131.1, 129.1, 129.0, 128.6, 127.3, 126.4, 123.4, 121.0, 118.5, 114.2, 55.5, 19.3. MS (ESI) 364 ( $\text{MH}^+$ ).


Example 113 – 2-(4-Chloro-2-nitro-phenylsulfanyl)-benzoic acid methyl ester (189JO09)



[0269] To a mixture of 5-chloro-2-nitrofluorobenzene (176 mg, 1 mmol) and methyl thiosalicylate (275  $\mu\text{L}$ , 2 mmol) in DMF (5 mL) was added  $\text{Cs}_2\text{CO}_3$  (652 mg, 2 mmol) and the resulting mixture was stirred at room temperature for 2 h. The mixture was diluted with  $\text{CH}_2\text{Cl}_2$ , washed with water, dried ( $\text{Na}_2\text{SO}_4$ ), concentrated and flash


chromatographed (SiO<sub>2</sub>, heptane:toluene, 1:10-1:4) to give 300 mg (92%) of the title compound (189JO09). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.15 (d, 1 H *J* = 2.4 Hz), 7.94 (m, 1 H), 7.53-7.46 (m, 3 H), 7.34 (dd, 1 H, 2.4, 8.6 Hz), 6.95 (d, 1 H, *J* = 8.8 Hz), 3.82 (s, 3 H).

Example 114 – 2-(2-Amino-4-chlorophenylsulfanyl)-benzoic acid methyl ester (189JO11)




**[0270]** To a mixture of 2-(4-chloro-2-nitro-phenylsulfanyl)-benzoic acid methyl ester (189JO09) (232 mg, 0.72 mmol) in EtOH (5 mL) was added SnCl<sub>2</sub>·2H<sub>2</sub>O (812 mg, 3.6 mmol) and the resulting mixture was stirred at 80°C for 2 h and then concentrated. The residue was treated with ice, and then Na<sub>2</sub>CO<sub>3</sub> was added until a pH of 10 was reached. EtOAc was added and the slurry was filtered through celite. The EtOAc-phase was washed with water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to give 149 mg (70%) of the title compound (189JO11). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.02 (dd, 1 H, *J* = 1.6, 7.8 Hz), 7.39 (d, 1 H, *J* = 8.2 Hz), 7.29 (m, 1 H), 7.15 (dt, 1 H, *J* = 1.2, 7.8 Hz), 6.87 (d, 1 H, *J* = 2.2 Hz), 6.80 (dd, 1 H, *J* = 2.2, 8.2 Hz), 6.76 (dd, 1 H, *J* = 1.2, 8.0 Hz), 3.96 (s, 3 H).


Example 115 – 8-Chloro-10*H*-dibenzo[*b,f*][1,4]thiazepin-11-one (189JO13)




**[0271]** A mixture of 2-(2-amino-4-chlorophenylsulfanyl)-benzoic acid methyl ester (189JO11) (149 mg, 0.51 mmol) and AlMe<sub>3</sub> (355 μL, 0.71 mmol, 2 M in toluene) in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) was stirred at ambient temperature for six days, and then water was added carefully. The mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub>, and was acidified with 2 M aqueous HCl. The organic phase was separated, dried (Na<sub>2</sub>SO<sub>4</sub>), concentrated and flash chromatographed (heptane:EtOAc, 5:1-3:1) to give 38 mg (29%) of the title compound (189JO13). MS (ESI) 262 (MH<sup>+</sup>).

Example 116 – 2-(Chloro-2-nitro-phenoxy)-benzoic acid methyl ester (189JO29A)


[0272]  $\text{Cs}_2\text{CO}_3$  (1.30 g, 4 mmol) was added to a mixture of 5-chloro-2-nitrofluorobenzene (352 mg, 2 mmol) and methyl 2-hydroxybenzoate (0.52 mL, 4 mmol) in DMF (6 mL) and the resulting mixture was stirred at room temperature for 2 h. The mixture was diluted with  $\text{CH}_2\text{Cl}_2$ , washed with water, dried ( $\text{Na}_2\text{SO}_4$ ), concentrated and flash chromatographed ( $\text{SiO}_2$ , heptane:EtOAc, 10:1-4:1) to give 505 mg (82%) of the title compound (189JO29A).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  8.02 (dd, 1 H,  $J = 1.8, 7.8$  Hz), 7.96 (d, 1 H,  $J = 1.9$  Hz), 7.59 (dt, 1 H,  $J = 2.0, 7.6$  Hz), 7.39 (dd, 1 H,  $J = 2.5, 9.0$  Hz), 7.24 (dt, 1 H,  $J = 1.2, 7.6$  Hz), 7.13 (dd, 1 H,  $J = 1.2, 8.0$  Hz), 6.74 (d, 1 H,  $J = 9.0$  Hz), 3.77 (s, 3 H).

Example 117 – 8-Chloro-10H-dibenzo[b,f][1,4]oxazepin-11-one (189JO29C)

[0273] Pd (catalytic amount, 5 % on carbon) was added to a solution of 2-(chloro-2-nitro-phenoxy)-benzoic acid methyl ester (189JO29A) (505 mg, 1.64 mmol) in EtOAc (20 mL) and the resulting mixture was hydrogenated ( $\text{H}_2$ , 1 atm.) for 48 h, then filtered through celite and concentrated. The residue was taken up in toluene (6 mL) and NaH (160 mg, 4.0 mmol, 60% in mineral oil) was added. The resulting mixture was stirred at  $80^\circ\text{C}$  over night, and then quenched by addition of saturated aqueous  $\text{NH}_4\text{Cl}$ -solution. The resulting mixture was diluted with EtOAc, washed with water, dried ( $\text{Na}_2\text{SO}_4$ ), concentrated and flash chromatographed ( $\text{SiO}_2$ , toluene:EtOAc, 4:1), which gave 171 mg (42%) of the title compound (189JO29C).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  8.12 (bs, 1 H), 7.95 (dd, 1 H,  $J = 1.8, 8.0$  Hz), 7.54 (dt, 1 H,  $J = 1.8, 8.0$  Hz), 7.29-7.19 (m, 3 H), 7.08 (dd, 1 H,  $J = 2.3, 8.6$  Hz), 7.04 (d, 1 H,  $J = 2.3$  Hz). MS (ESI) 246 ( $\text{MH}^+$ ).

Example 118 – 3-Chloro-5,11-dihydro-dibenzo[*b,e*]azepin-6-one (189JO59)

[0274] To a mixture of 5-chloro-2-methylphenyl isocyanate (100  $\mu$ L, 0.73 mmol) in  $\text{CCl}_4$  (2 mL) was added sulfuryl chloride (118  $\mu$ L, 0.88 mmol) and 2,2'-azobis(isobutyronitrile) (catalytic amount) and the resulting mixture was refluxed for 20h. The mixture was allowed to obtain room temperature, then diluted with  $\text{CH}_2\text{Cl}_2$ , washed with saturated aqueous  $\text{NaHCO}_3$ -solution, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated. The mixture was taken up in benzene (2 mL) and a mixture of  $\text{AlCl}_3$  (160 mg, 1.2 mmol) in benzene (1 mL) was added. The resulting mixture was stirred at  $80^\circ$  for 4h, and then allowed to obtain room temperature. The mixture was filtered through a short column ( $\text{SiO}_2$ , heptane:EtOAc, 1:1) to give 25 mg (14%) of the title compound (189JO59).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  8.18 (bs, 1 H), 7.92 (dd, 1 H,  $J = 1.2, 7.8$  Hz), 7.46 (dt 1 H,  $J = 1.4, 7.4$  Hz), 7.34 (dt, 1 H,  $J = 1.2, 7.4$  Hz), 7.23 (m, 2 H), 7.07 (m, 2 H), 3.92 (s 2 H). MS (ESI) 244 ( $\text{MH}^+$ )

Example 119 – 8-Bromo-10*H*-dibenzo[*b,f*][1,4]oxazepin-11-one (189JO56)

[0275] A mixture of a methyl 2-hydroxybenzoate (1.0 mL, 10.0 mmol), 5-bromo-2-fluoronitrobenzene (0.62 mL, 5.0 mmol) and  $\text{Cs}_2\text{CO}_3$  (3.3 g, 10.0 mol) in DMF (12 mL) was stirred at  $40^\circ\text{C}$  for 2h. The mixture was diluted with EtOAc and washed with 2 M aqueous  $\text{NaOH}$ -solution. To the EtOAc-phase was added EtOH,  $\text{H}_2\text{O}$ ,  $\text{K}_2\text{CO}_3$  (2.8 g, 20 mmol) and  $\text{Na}_2\text{S}_2\text{O}_4$  (3.5 g, 20 mmol) and the resulting mixture was stirred vigorously for 1 h. The aqueous phase was removed and the organic phase was washed with 1 M aqueous  $\text{NaOH}$ -solution and then concentrated. The residue was taken up in DMF (1 mL) and then toluene (4 mL) and  $\text{NaH}$  (60 mg, 1.5 mmol, 60% in mineral oil) were added and the resulting mixture was stirred at  $80^\circ\text{C}$  over night, then quenched by addition of saturated aqueous  $\text{NH}_4\text{Cl}$ -solution. The resulting mixture was diluted with EtOAc, washed with 2 M aqueous  $\text{NaOH}$ -solution, dried ( $\text{Na}_2\text{SO}_4$ ), concentrated, filtered through a short  $\text{SiO}_2$ -




column, concentrated and crystallised from heptane:EtoAc to give 130 mg of the title compound (189JO56). MS (ESI) 290 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 120 – General procedure 8 (GP8)


[0276] A BOC-protected diamine (1.8 eq.) was added to 8-chloro-11-methylsulfanyl-5*H*-dibenzo[*b,e*][1,4]diazepine(166JO50) (purity 50%, 1 eq.) in pyridine. The resulting mixture was heated in a capped tube at 110°C for 66 h. The mixture was concentrated and then diluted with  $CH_2Cl_2$ :trifluoroacetic acid (2:1-ratio). The resulting mixture was stirred at ambient temperature over night, and then concentrated. The residue was taken up in  $CH_2Cl_2$  and washed with saturated aqueous  $NaHCO_3$ -solution. The organic phase was applied onto a SCX-2 ion exchange column. The column was washed with MeOH, and then the product was eluted with  $NH_3$  (7 N in MeOH), concentrated and purified on HPLC.

Example 121 – (8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-(S)-1-pyrrolidin-2-yl-methyl-amine (166JO51)



[0277] 8-Chloro-11-methylsulfanyl-5*H*-dibenzo[*b,e*][1,4]diazepine (166JO50) (50 mg, 0.11 mmol) and (S)-(2-aminomethyl)-1-*N*-(*tert*-butoxycarbonylamino)-pyrrolidine (39 mg, 0.2 mmol) were reacted according to GP8 to give 3.0 mg of the title compound (166JO51). MS (ESI) 327 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/92.


Example 122 – 1-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-piperidine-4-yl-amine (166JO55)



[0278] 8-Chloro-11-methylsulfanyl-5*H*-dibenzo[*b,e*][1,4]diazepine (166JO50) (50 mg, 0.11 mmol) and 4-(*tert*-butoxycarbonylamino)-aminopiperidine (39 mg, 0.2 mmol)


were reacted according to GP8 to give 6.5 mg of the title compound (166JO55). MS (ESI) 327 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/99.

Example 123 – 1-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-pyrrolidin-3-yl-amine (166JO64)




**[0279]** 8-Chloro-11-methylsulfanyl-5*H*-dibenzo[*b,e*][1,4]diazepine (166JO50) (100 mg, 0.22 mmol) and 3-(*tert*-butoxycarbonylamino)pyrrolidine (73 mg, 0.4 mmol) were reacted according to GP8 to give 8.1 mg of the title compound (166JO64). MS (ESI) 313 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/94.

Example 124 – (8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-(R)-1-pyrrolidin-2-yl-methyl-amine (166JO70)




**[0280]** 8-Chloro-11-methylsulfanyl-5*H*-dibenzo[*b,e*][1,4]diazepine (166JO50) (100 mg, 0.22 mmol) and (R)-(2-aminomethyl)-1-*N*-(*tert*-butoxycarbonylamino)-pyrrolidine (78 mg, 0.4 mmol) were reacted according to GP8 to give 7.6 mg of the title compound (166JO70). MS (ESI) 327 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/90.

Example 125 – (8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-pyrrolidin-3-yl-amine (166JO74)




[0281] 8-Chloro-11-methylsulfanyl-5*H*-dibenzo[*b,e*][1,4]diazepine (166JO50) (100 mg, 0.22 mmol) and 3-amino-1-*N*-(*tert*-butoxycarbonylamino)pyrrolidine (73 mg, 0.4 mmol) were reacted according to GP8 to give 7.7 mg of the title compound (166JO74). MS (ESI) 313 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/90.

Example 126 – 8-Chloro-11-(2,5-diaza-bicyclo[2.2.1]hept-2-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine (166JO39-2)



[0282] 8-Chloro-11-methylsulfanyl-5*H*-dibenzo[*b,e*][1,4]diazepine (166JO50) (50 mg, 0.11 mmol) and *N*-(*tert*-butoxycarbonylamino)-2,5-diazabicyclo[2.2.1]heptane (34 mg, 0.2 mmol) were reacted according to GP8 to give 15 mg of the title compound (166JO39-2). MS (ESI) 324 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 93/100.

Example 127 – Acetidin-3-yl-(8-chloro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)amine (189JO65)



[0283] To 8,5-Dichloro-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE64) (30 mg, 0.11 mmol) in dioxane (2.0 mL) was added 3-amino-azetidine-1-carboxylic acid *tert*-butyl ester (59 mg, 0.34 mmol) and  $CS_2CO_3$  (74 mg, 0.23 mmol) and the resulting mixture was heated in capped tube using microwave assisted heating (170°C, 40 minutes). The mixture was diluted with EtOAc, washed with water, dried ( $Na_2SO_4$ ) and concentrated. The residue was taken up in  $CH_2Cl_2$  (2 mL) and trifluoroacetic acid (1 mL) was added. The resulting mixture was stirred at ambient temperature over night, and then concentrated. The residue was taken up in  $CH_2Cl_2$  and washed with saturated aqueous  $NaHCO_3$ -solution. The organic phase was applied onto a SCX-2 ion exchange column. The column was washed with MeOH, and then the product was eluted with  $NH_3$  (7 N in MeOH), concentrated, and

purified by HPLC to give 16 mg of the title compound (189JO65). MS (ESI) 299 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 97/90.


Example 128 – General procedure 9 (GP9)

[0284] A mixture of a 3-aminomethyl ester (1 eq.), 5-bromo-2-fluoronitrobenzene (1 eq.) and  $K_2CO_3$  (4 eq.) in DMF was heated to 60°C for 1 hour, and then allowed to obtain room temperature. The mixture was diluted with  $CH_2Cl_2$  and washed with saturated aqueous  $NH_4Cl$ -solution, dried ( $Na_2SO_4$ ) and concentrated. The residue was taken up in EtOH and a mixture of  $K_2CO_3$  (5 eq.) and  $Na_2S_2O_4$  (5 eq.) in water was added and the resulting mixture was stirred vigorously for 1 h. The aqueous phase was extracted with EtOAc (3 x) and the combined organic phases were dried ( $Na_2SO_4$ ) and concentrated.

[0285] The residue was taken up in  $CH_3CN$ ,  $H_2SO_4$  (10 vol-%, 98%) was added, and the resulting mixture was stirred at 80°C for 1 h. The mixture was diluted with  $CH_2Cl_2$ , washed with saturated aqueous  $NaHCO_3$ -solution, dried ( $Na_2SO_4$ ), concentrated, flash chromatographed ( $SiO_2$ , heptane:EtOAc-system), and concentrated to give intermediate lactam.

[0286] The residue was taken up in dioxane and added to a mixture of  $TiCl_4$  (1.1 eq., 1 M in toluene) and piperazine (5 eq.) in dioxane at 50°C. The resulting mixture was stirred at 100°C over night, and then allowed to obtain room temperature. To the mixture was added aqueous HCl (2 M) until acidic solution and then the aqueous phase was extracted with EtOAc (2 x). To the aqueous phase was added aqueous NaOH (2 M) until basic solution and the resulting suspension was extracted with EtOAc (3 x). The combined organic phases were concentrated and flash chromatographed ( $SiO_2$ ,  $CH_2Cl_2$ :MeOH,  $NH_3$ (7N in MeOH))-system.


Example 129 – 7-Bromo-4-(piperazin-1-yl)-2,3-dihydro-1H-benzo[b][1,4]diazepine (166JO47)



[0287] 5-Bromo-2-fluoronitrobenzene (440 mg, 2.0 mmol) and methyl 3-amino propionate hydrochloride (920 mg, 3.0 mmol) were reacted according to GP9 to give 4.0


mg of the title compound (166JO47). MS (ESI) 309 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 130 – 7-Bromo-2-methyl-(piperazin-1-yl)-2,3-dihydro-1H-benzo[*b*][1,4]diazepine (166JO95)




[0288] 5-Bromo-2-fluoronitrobenzene (440 mg, 2.0 mmol) and methyl 3-amino buturate (787 mg, 3.0 mmol) were reacted according to GP9 to give 12 mg of the title compound (166JO95). MS (ESI) 323 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 100/100.

Example 131 – 7-Bromo-2-phenyl-4-(piperazine-1-yl)-2,3-dihydro-1H-benzo[*b*][1,4]diazepine (189JO20)



[0289] 5-Bromo-2-fluoronitrobenzene (440 mg, 2.0 mmol) and ethyl 3-amino-3-phenylpropionate hydrochloride (394 mg, 1.5 mmol) were reacted according to GP9 to give 9.8 mg of the title compound (189JO20). MS (ESI) 385 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 97/88.

Example 132 – 7-Bromo-10-(piperazin-1-yl)-1,2,3,3a,4,10a-hexahydro-benzo[*b*]cyclopenta[*e*][1,4]diazepine (166JO46)




[0290] 5-Bromo-2-fluoronitrobenzene (110 mg, 0.5 mmol) and cis-2-amino-1-cyclopentanecarboxylic acid hydrochloride (138 mg, 0.75 mmol) were reacted according to GP2 to give 3.0 mg of the title compound (166JO46). MS (ESI) 349 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 99/88.

Example 133 – General Procedure 10 (GP10)


[0291] A zinc reagent (0.4 mmol) was added at room temperature to a solution of 8,5-Dichloro-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE64) (53mg, 0.2 mmol) and  $PdCl_2(PPh_3)_2$  (9 mg, 0.02 mmol) in dry THF (1 ml). The reaction was shaken until complete conversion (1-16h, TLC) and then quenched by the addition of aqueous saturated  $NH_4Cl$ . The resulting mixture was extracted twice with ether and the combined ethereal phases were washed with brine and dried over  $Na_2SO_4$ . Filtration followed by concentration at reduced pressure of the organic phase gave a crude product, which was purified using column chromatography (heptane:EtOAc-system).

Example 134 – 8-Chloro-11-(4-fluorobenzyl)-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE59)



[0292] 4-Fluorobenzylzinc chloride (0.8 ml, 0.5 M in THF, 0.4 mmol) and 8,5-dichloro-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE64) (53 mg, 0.2 mmol) were reacted according GP10 to give 52 mg of the title compound (160FE59). MS (ESI) 337 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 90/90.

Example 134 – 8-Chloro-11-(4-fluorophenyl)-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE70)




[0293] 4-Fluorophenylzinc chloride (0.5 ml, 0.5 M in THF, 0.4 mmol) and 8,5-dichloro-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE64) (26 mg, 0.1 mmol) were reacted according GP10 to give 23 mg of the title compound (160FE70). MS (ESI) 323 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 98/100.

Example 135 – General procedure 11 (GP11)


[0294] Aqueous Na<sub>2</sub>CO<sub>3</sub> (1 ml, 1M) was added at room temperature to a solution of the 8,5-dichloro-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE64) (53mg, 0.2 mmol) (26mg, 0.1 mmol), Pd(PPh<sub>3</sub>)<sub>4</sub> (10 mg), and the appropriate boronic acid reagent (0.12 mmol) in dioxane (3 ml). The mixture was then shaken at 80°C until complete conversion of the imidoyl chloride (TLC). The temperature was decreased and ether and H<sub>2</sub>O were added to the reaction mixture. The ether phase was washed with brine and dried over Na<sub>2</sub>SO<sub>4</sub>. Filtration followed by concentration at reduced pressure of the organic phase gave a crude product, which was purified using column chromatography (heptane:EtOAc-system).

Example 136 – 8-Chloro-11-(4-nonylphenyl)-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE63)




[0295] 4-Nonylphenylboronic acid (30 mg, 0.12 mmol) and 8,5-dichloro-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE64) (26 mg, 0.1 mmol) were reacted according GP11 to give 25 mg of the title compound (160FE63). MS (ESI) 431 (MH<sup>+</sup>). Purity for MH<sup>+</sup> (UV/MS) 85/85.

Example 137 – 8-Chloro-11-(pyridin-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE69A)



[0296] 4 pyridyl-4-boronic acid (14 mg, 0.12 mmol) and 8,5-dichloro-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE64) (26 mg, 0.1 mmol) were reacted according GP11 to give 9.3 mg of the title compound (160FE69A). MS (ESI) 306 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 98/95.

Example 138 – 8-Chloro-11-(1*H*-pyrazol-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE59)



[0297] 4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole (22 mg, 0.12 mmol) and 8,5-dichloro-5*H*-dibenzo[*b,e*][1,4]diazepine (160FE64) (26 mg, 0.1 mmol) were reacted according GP11 to give 8.7 mg of the title compound (160FE69B). MS (ESI) 295 ( $MH^+$ ). Purity for  $MH^+$  (UV/MS) 95/100.

Example 139 – Functional Screen

[0298] Several compounds disclosed herein were evaluated for activity at muscarinic receptors using Receptor Selection and Amplification Technology (R-SAT) as described in U.S. Patent No. 5,707,798, the disclosure of which is incorporated herein by reference in its entirety. The efficacy (eff) and potency (expressed as  $pEC_{50}$ ) of these compounds are presented in Table 1 at M1, M2, and M3 receptors.



Table 1.

Name	M1		M2		M3	
	eff	pEC <sub>50</sub>	eff	pEC <sub>50</sub>	eff	pEC <sub>50</sub>
2,7-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO85F1)	-2		-14		-	
2-Chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO85F6)	-5		8		-	
2,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO85F2)	1		24		-	
8-Bromo-2-chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO85F3)	10		29	5.5	-	
2-Chloro-11-(piperazin-1-yl)-8-trifluoromethyl-5H-dibenzo[b,e][1,4]diazepine (166JO85F7)	10		19		-	
6-Chloro-11-(piperazin-1-yl)-8-trifluoromethyl-5H-dibenzo[b,e][1,4]diazepine (189JO77B)	-		-		-	
7-Chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE35B)	48	6.6	53	5.4	-	
8-Bromo-1-chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE36A)	66	6.9	105	5.6	4	
8-Bromo-2-methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE40C)	8		44	5.4	-	
4,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE41A)	78	7.2	118	6.0	44	6.8
8-Chloro-2-methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE41B)	14		28	5.9	-	
8-Chloro-2-fluoro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE42A-F3)	32	6.8	121	5.5	-	
3,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE42B-F4)	34	6.9	58	5.8	-	
2-Bromo-8-chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE43A-F6)	6		19		-	

3,7-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE58D1)	15	19	-	-
8-Bromo-3-chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE58D3)	31	34	6.5	-
3-Chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE58D6)	16	27	5.7	-
3-Chloro-11-(piperazin-1-yl)-8-trifluoromethyl-5H-dibenzo[b,e][1,4]diazepine (160FE58D7)	11	-	-	-
7-Chloro-2-methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE58E1)	-3	5	-	-
2-Methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE58E6)	1	9	-	-
2-Methyl-11-(piperazin-1-yl)-8-trifluoromethyl-5H-dibenzo[b,e][1,4]diazepine (160FE58E7)	15	14	-	-
8-Chloro-4-methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE74C)	92	162	6.0	16
1,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (203FE03)	-	84	5.7	-
8-Bromo-5-methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO32)	92	75	5.6	-
7,8-Dichloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO28)	32	8	-	-
11-(Piperazin-1-yl)-8-trifluoromethyl-5H-dibenzo[b,e][1,4]diazepine (166JO23)	53	131	5.5	-
11-(Piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE19A)	38	70	5.6	-
8-Fluoro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE19C)	32	95	5.7	-
11-(Piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine-8-carbonitrile (160FE19D)	49	106	6.0	-
8-Bromo-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE19E)	63	121	6.3	13
8-Methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE19F)	49	98	6.0	-

3-Fluoro-6-piperazin-1-yl-1,1,1H-dibenzo[b,e]azepine (160FE02)	23	55	6.1	-	
2-(Trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE13A)	-8	9		-	
2-(Trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]oxazepine (160FE13B)	-11	4		-	
8-Chloro-2-(trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE13C)	-2	16		-	
8-(Trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE13D)	19	62	5.6	-	
11-(Piperazin-1-yl)-dibenzo[b,f][1,4]thiazepin (160FE17A)	4	41	6.5	-	
11-(Piperazin-1-yl)-2,3-dihydro-1,4-benzodioxino[6,7-b][1,4]benzothiazepin (160FE17B)	6	28			
8-Chloro-11-[1,4]diazepam-1-yl-5H-dibenzo[b,e][1,4]diazepine (160FE16A)	37	82	6.2	-	
N <sup>7</sup> -(8-Chloro-5H-dibenzo[b,e][1,4]diazepine-11-yl)-N,N-dimethylethane-1,2-diamine (160FE16D)	4	11		-	
N <sup>7</sup> -(8-Chloro-5H-dibenzo[b,e][1,4]diazepine-11-yl)-N,N-diethyl-ethane-1,2-diamine (160FE16E)	6	9		-	
8-Chloro-11-(4-methyl-[1,4]diazepam-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE16F)	18	50	5.9	-	
8-Chloro-2-methoxy-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE20A)	7	7		-	
N <sup>7</sup> -(5H-dibenzo[b,e][1,4]diazepine-11-yl)-N,N-dimethyl-ethane-1,2-diamine (160FE20B)	10	20		-	
11-[1,4]Diazepam-1-yl-5H-dibenzo[b,e][1,4]diazepine (160FE20C)	18	25		-	
N <sup>7</sup> -(8-Fluoro-5H-dibenzo[b,e][1,4]diazepine-11-yl)-N,N-dimethyl-ethane-1,2-diamine (160FE20D)	7	21		-	
8-Fluoro-11-[1,4]diazepam-1-yl-5H-dibenzo[b,e][1,4]diazepine (160FE16A)	25	67	6.6	-	
N <sup>7</sup> -(8-Chloro-5H-dibenzo[b,e][1,4]diazepine-11-yl)-N-methyl-ethane-1,2-	5	8		-	

diamine (160FE22)									
8-Chloro-11-(trans-2,5-dimethyl-piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE33A)	8				24			-	
8-Chloro-11-(3,5-dimethyl-piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE33B)	18				97		5.6	-	
8-Chloro-11-(3-methyl-piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE38)	28		7.0		159		5.8	19	
8-Chloro-11-(3-phenyl-piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE45)	5				23			-	
8-Chloro-5-methyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (189JO25A)	65		7.1		138		5.9	14	
8-Chloro-5-benzyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE46-PIPN)	41		6.3		16			-	
8-Iodo-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO38)	75		7.2		187		6.0	-	
2-Iodo-8-chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO54)	3				10			-	
8-Phenyl-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (189JO53)	-				104		5.7	-	
8-Chloro-11-(piperidin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO69A)	47		5.8		9			-	
8-Chloro-11-(morpholin-4-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO69B)	11				6			-	
5-Allyl-8-chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO68)	12				46		5.8	-	
6-Chloro-11-(piperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine (189JO68)	-				71		5.4	-	
8-Chloro-5-piperazin-1-yl-11H-benzo[b]pyrido[2,3-e][1,4]diazepine (166JO63)	51		5.5		7			-	
2-Chloro-10-piperazin-1-yl-5H-dibenzo[b,f]azepin (189JO39)	2				3			-	
8-Chloro-11-(piperazin-1-yl)-dibenzo[b,f][1,4]thiazepine (189JO16)	11				52		5.9	-	
8-Chloro-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO31)	52		7.0		58		6.0	-	
8-Chloro-11-(4-methyl-piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO47)	13				46		5.8	-	


3-Chloro-6-piperazin-1-yl-11H-dibenzo[b,e]azepine (189JO60)	-	113	5.9	-
8-Bromo-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO48A)	-	70	5.9	-
11-(Piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO48B)	-	58	5.7	-
7-Chloro-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO50A)	-	53	5.9	-
8-Chloro-3-methoxy-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO50B)	-	19		-
8-Bromo-3-methoxy-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO50D)	-	21		-
3-Methoxy-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO50E)	-	45	5.6	-
7-Chloro-3-methoxy-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO50F)	-	44	6.0	-
8-Chloro-4-methyl-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO50H)	-	71	6.2	-
8-Bromo-4-methyl-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO51A)	-	48	5.8	-
4-Methyl-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO51B)	-	22		-
2-Bromo-8-chloro-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO51D)	-	13		-
2,8-Dibromo-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO51E)	-	6		-
2-Bromo-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO51F)	-	9		-
2-Bromo-7-chloro-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO51G)	-	15		-
11-(Piperazin-1-yl)-8-trifluoromethyl-dibenzo[b,f][1,4]oxazepine (189JO54A)	-	58	5.9	-
4-Methyl-11-(piperazin-1-yl)-8-trifluoromethyl-dibenzo[b,f][1,4]oxazepine (189JO54C)	-	75	6.1	-
8-Fluoro-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO54E)	-	48	5.6	-
8-Fluoro-3-methoxy-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO54F)	-	49	5.6	-
8-Fluoro-4-methyl-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine	-	50	5.8	-

(189JO54G)									
2-Bromo-8-fluoro-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO54H)	-				21			-	
8-Methyl-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO58A)	-				76		5.5	-	
3-Methoxy-8-methyl-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO58B)	-				45		5.5	-	
4,8-Dimethyl-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO58C)	-				46		6.1	-	
3-Methoxy-11-(piperazin-1-yl)-8-trifluoromethyl-dibenzo[b,f][1,4]oxazepine (189JO62A)	-				17			-	
2-Bromo-11-(piperazin-1-yl)-8-trifluoromethyl-dibenzo[b,f][1,4]oxazepine (189JO62B)	-				11			-	
6-Chloro-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO69)	-				-			-	
2-Bromo-8-methyl-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO63A)	-				17			-	
7-Chloro-4-methyl-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO63B)	-				28		6.4	-	
8-Phenyl-11-(piperazin-1-yl)-dibenzo[b,f][1,4]oxazepine (189JO64)	-				33		5.9	-	
8-Chloro-11-(piperidin-4-yl)-5H-dibenzo[b,e][1,4]diazepine(160FE67A)	57	6.2			31		5.7	-	
5-Benzyl-8-chloro-11-(piperidin-4-yl)-5H-dibenzo[b,e][1,4]diazepine(160FE67B)	12				5			-	
(8-Chloro-5H-dibenzo[b,e][1,4]diazepin-11-yl)-(S)-1-pyrrolidin-2-yl-methyl-amine (166JO51)	77	6.3			23			-	
1-(8-Chloro-5H-dibenzo[b,e][1,4]diazepin-11-yl)-piperidine-4-yl-amine (166JO55)	6				95		5.6	-	
1-(8-Chloro-5H-dibenzo[b,e][1,4]diazepin-11-yl)-pyrrolidin-3-yl-amine (166JO64)	8				36		5.5	-	
(8-Chloro-5H-dibenzo[b,e][1,4]diazepin-11-yl)-(R)-1-pyrrolidin-2-yl-methyl-amine (166JO70)	4				5			-	
(8-Chloro-5H-dibenzo[b,e][1,4]diazepin-11-yl)-pyrrolidin-3-yl-amine (166JO74)	10				18			-	

8-Chloro-11-(2,5-diaza-bicyclo[2.2.1]hept-2-yl)-5H-dibenzo[b,e][1,4]diazepine (166JO39-2)	34	6.6	49	5.6	-	
Acetidin-3-yl-(8-chloro-5H-dibenzo[b,e][1,4]diazepine-11-yl)amine (189JO65)	-		17		-	
7-Bromo-4-(piperazin-1-yl)-2,3-dihydro-1H-benzo[b][1,4]diazepine (166JO47)	56	5.6	3		-	
7-Bromo-2-methyl-(piperazin-1-yl)-2,3-dihydro-1H-benzo[b][1,4]diazepine (166JO95)	16		23		-	
7-Bromo-2-phenyl-4-(piperazine-1-yl)-2,3-dihydro-1H-benzo[b][1,4]diazepine (189JO20)	3		19		-	
7-Bromo-10-(piperazin-1-yl)-1,2,3,3a,4,10a-hexahydro-benzo[b]cyclopenta[e][1,4]diazepine (166JO46)	50	5.8	12		-	
8-Chloro-11-(4-fluorobenzyl)-5H-dibenzo[b,e][1,4]diazepine (160FE59)	3		7		-	
8-Chloro-11-(4-fluorophenyl)-5H-dibenzo[b,e][1,4]diazepine (160FE70)	13		4		-	
8-Chloro-11-(4-nonylphenyl)-5H-dibenzo[b,e][1,4]diazepine (160FE63)	1		8		-	
8-Chloro-11-(pyridin-4-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE69A)	6		6		-	
8-Chloro-11-(1H-pyrazol-4-yl)-5H-dibenzo[b,e][1,4]diazepine (160FE59)	9		3		-	


WHAT IS CLAIMED IS:

1. A compound of Formula I, II, or XV:



or a pharmaceutically acceptable salt, ester, amide, or prodrug thereof, wherein:

A is selected from the group consisting of



X is nitrogen, CH, or CH<sub>2</sub>;

X' is C or CH, wherein when X' is C, there is a double bond between X and X' and wherein when X' is CH, there is a single bond between X and X';

each Y is separately selected from the group consisting of nitrogen, oxygen, or CH;

each W is separately selected from the group consisting of nitrogen, CH, oxygen, or sulfur;

each n is separately selected from the group consisting of 0, 1, 2, 3, and 4;

m is selected from the group consisting of 1, 2, and 3;



each  $R_1$  is separately absent or is separately selected from the group consisting of hydrogen, halogen, amine, optionally substituted  $C_{1-20}$  alkyl, optionally substituted  $C_{3-8}$  cycloalkyl, optionally substituted  $C_{2-20}$  alkenyl, optionally substituted  $C_{2-20}$  alkynyl, optionally substituted  $C_{1-20}$  -alkoxyalkyl, and optionally substituted aryl and arylalkyl;

L is absent or is selected from the group consisting of  $-NH(CH_2)_n-$  and  $-(CH_2)_n-$ ;

a, b, c, and d are each separately selected from the group consisting of carbon, nitrogen, oxygen, and sulfur, or each is separately absent,

provided that at least three of a, b, c, or d are present,

provided that at least one of a, b, c, or d is carbon, and

provided that no two adjacent a, b, c, or d are both oxygen or both sulfur;

e, f, g, and h are each separately selected from the group consisting of carbon, nitrogen, oxygen, and sulfur, or each is separately absent,

provided that at least three of e, f, g, or h are present,

provided that at least one of e, f, g, or h is carbon, and

provided that no two adjacent e, f, g, or h are both oxygen or both sulfur;

$R_2$ ,  $R_3$ ,  $R_4$ , and  $R_5$ , are each separately selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, optionally substituted  $C_{1-6}$  alkyloxy, optionally substituted  $C_{2-6}$  alkenyl, optionally substituted  $C_{2-6}$  alkynyl, optionally substituted  $C_{1-6}$ -alkoxyalkyl, optionally substituted  $C_{1-6}$  alkylthio, perhaloalkyl, CN,  $COR_{10}$ ,  $CONHR_{10}$ ,  $NHCONHR_{10}$ ,  $SO_2NHR_{10}$ ,  $SO_2R_{10}$ ,  $OSO_2R_{10}$ , heteroalkyl,  $NO_2$ ,  $NHCOR_{10}$ ,

or  $R_2$  and  $R_3$ , or  $R_3$  and  $R_4$ , or  $R_4$  and  $R_5$  taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety;

$R_6$ ,  $R_7$ ,  $R_8$ , and  $R_9$ , are each separately selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, optionally substituted  $C_{1-6}$  alkyloxy, optionally substituted  $C_{2-6}$  alkenyl, optionally substituted  $C_{2-6}$  alkynyl, optionally substituted  $C_{1-6}$ -alkoxyalkyl, optionally substituted  $C_{1-6}$  alkylthio, perhaloalkyl, CN,  $COR_{10}$ ,  $CONHR_{10}$ ,  $NHCONHR_{10}$ ,  $SO_2NHR_{10}$ ,  $SO_2R_{10}$ ,  $OSO_2R_{10}$ , heteroalkyl,  $NO_2$ ,  $NHCOR_{10}$ ,

or R<sub>6</sub> and R<sub>7</sub>, or R<sub>7</sub> and R<sub>8</sub>, or R<sub>8</sub> and R<sub>9</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety;

Z is selected from the group consisting of NR<sub>11</sub>, oxygen, sulfur, and CH<sub>2</sub>;

R<sub>10</sub> is selected from the group consisting of hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl optionally substituted aryl, optionally substituted arylalkyl, and perhaloalkyl; and

R<sub>11</sub> is selected from the group consisting of hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, and optionally substituted arylalkyl;


R<sub>12</sub> and R<sub>13</sub> are separately selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>,

or R<sub>12</sub> and R<sub>13</sub>, taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety;


any bond represented by a dashed and solid line represents a bond selected from the group consisting of a carbon-carbon single bond and a carbon-carbon double bond;

provided that the compound of Formulae I or XV are not clozapine or N-desmethylozapine.



2. The compound of claim 1, wherein said compound has a structure set forth in Formulas III or IV.



3. The compound of claim 1, wherein said compound is selected from the group consisting of:



4. The compound of claim 3, wherein said compound is selected from the group consisting of:



5. The compound of claim 1, wherein none of a, b, c, or d is absent.
6. The compound of claim 1, wherein none of e, f, g, or h is absent.
7. The compound of claim 1, wherein a, b, c, and d are carbon.
8. The compound of claim 1, wherein e, f, g, and h are carbon.
9. The compound of claim 1, wherein  $R_2$  is selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, and optionally substituted  $C_{1-6}$  alkyloxy.
10. The compound of claim 9, wherein said alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl.
11. The compound of claim 9, wherein said alkyloxy is selected from the group consisting of methoxy, ethoxy, propoxy, isopropoxy, butoxy, sec-butoxy, and tert-butoxy.
12. The compound of claim 9, wherein said halogen is selected from the group consisting of fluoro, chloro, and bromo.
13. The compound of claim 1, wherein  $R_2$  is selected from the group consisting of hydrogen, methyl, methoxy, and chloro.
14. The compound of claim 1, wherein  $R_3$  is selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, optionally substituted  $C_{1-6}$  alkyloxy, and  $NO_2$ .

15. The compound of claim 14, wherein said alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl.
16. The compound of claim 14, wherein said alkyloxy is selected from the group consisting of methoxy, ethoxy, propoxy, isopropoxy, butoxy, sec-butoxy, and tert-butoxy.
17. The compound of claim 14, wherein said halogen is selected from the group consisting of chloro, bromo, and iodo.
18. The compound of claim 1, wherein  $R_3$  is selected from the group consisting of hydrogen, methyl, methoxy, chloro, bromo, iodo, and  $\text{NO}_2$ .
19. The compound of claim 1, wherein  $R_4$  is selected from the group consisting of hydrogen, halogen, optionally substituted  $\text{C}_{1-6}$  alkyl, perhaloalkyl,  $\text{SO}_2\text{R}_{10}$ , and  $\text{NO}_2$ .
20. The compound of claim 19, wherein said alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl.
21. The compound of claim 19, wherein said perhaloalkyl is perfluoroalkyl.
22. The compound of claim 21, wherein said perfluoroalkyl is trifluoromethyl.
23. The compound of claim 19, wherein said halogen is selected from the group consisting of fluoro, chloro, and bromo.
24. The compound of claim 19, wherein  $\text{R}_{10}$  is hydrogen or optionally substituted  $\text{C}_{1-6}$  alkyl.
25. The compound of claim 24, wherein said alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl.
26. The compound of claim 1, wherein  $R_4$  is selected from the group consisting of hydrogen, methyl, fluoro, chloro, bromo, trifluoromethyl,  $\text{SO}_2\text{CH}_3$ , and  $\text{NO}_2$ .
27. The compound of claim 1, wherein  $R_5$  is selected from the group consisting of hydrogen, halogen, and optionally substituted  $\text{C}_{1-6}$  alkyl.
28. The compound of claim 27, wherein said alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl.
29. The compound of claim 27, wherein said halogen is selected from the group consisting of fluoro, chloro, and bromo.
30. The compound of claim 1, wherein  $R_5$  is hydrogen or chloro.
31. The compound of claim 1, wherein  $R_6$  is hydrogen or optionally substituted  $\text{C}_{1-6}$  alkyl.
32. The compound of claim 1, wherein  $R_6$  is hydrogen.

33. The compound of claim 1, wherein  $R_7$  is selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, perhaloalkyl, CN,  $SO_2R_{10}$ , and  $NO_2$ .
34. The compound of claim 33, wherein said alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl.
35. The compound of claim 33, wherein said halogen is selected from the group consisting of fluoro, chloro, and bromo.
36. The compound of claim 33, wherein said perhaloalkyl is perfluoroalkyl.
37. The compound of claim 36, wherein said perfluoroalkyl is trifluoromethyl.
38. The compound of claim 33, wherein  $R_{10}$  is hydrogen or optionally substituted  $C_{1-6}$  alkyl.
39. The compound of claim 38, wherein said alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl.
40. The compound of claim 1, wherein  $R_7$  is selected from the group consisting of hydrogen, methyl, chloro, trifluoromethyl,  $SO_2CH_3$ , CN, and  $NO_2$ .
41. The compound of claim 1, wherein  $R_8$  is selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl.
42. The compound of claim 41, wherein said alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl.
43. The compound of claim 41, wherein said halogen is selected from the group consisting of fluoro, chloro, and bromo.
44. The compound of claim 1, wherein  $R_8$  is selected from the group consisting of hydrogen, chloro, and bromo.
45. The compound of claim 1, wherein  $R_9$  is selected from the group consisting of hydrogen, halogen, optionally substituted  $C_{1-6}$  alkyl, and perhaloalkyl.
46. The compound of claim 45, wherein said alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl.
47. The compound of claim 45, wherein said halogen is selected from the group consisting of fluoro, chloro, and bromo.
48. The compound of claim 45, wherein said perhaloalkyl is perfluoroalkyl.
49. The compound of claim 48, wherein said perfluoroalkyl is trifluoromethyl.
50. The compound of claim 1, wherein  $R_9$  is selected from the group consisting of hydrogen, chloro, methyl, and trifluoromethyl.

51. The compound of claim 1, wherein  $R_1$  is selected from the group consisting of hydrogen, optionally substituted  $C_{1-6}$  alkyl, and optionally substituted aryl.

52. The compound of claim 51, wherein said alkyl is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, and tert-butyl.


53. The compound of claim 1, wherein  $R_1$  is hydrogen.

54. The compound of claim 1, wherein X is nitrogen.

55. The compound of claim 1, wherein Y is NH.

56. The compound of claim 1, wherein L is absent or is selected from the group consisting of  $-NHCH_2-$ ,  $-NH-$ , and  $-CH_2-$ .

57. The compound of claim 1, wherein A is selected from the group consisting of:



and wherein n is selected from the group consisting of 0, 1, and 2.

58. The compound of claim 1 selected from the group consisting of:

- 2,7-Dichloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 2-Chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 2,8-Dichloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 8-Bromo-2-chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 2-Chloro-11-(piperazin-1-yl)-8-trifluoromethyl-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 6-Chloro-11-(piperazin-1-yl)-8-trifluoromethyl-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 7-Chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 8-Bromo-1-chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 8-Bromo-2-methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 4,8-Dichloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 8-Chloro-2-methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 8-Chloro-2-fluoro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 3,8-Dichloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 2-Bromo-8-chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,
- 3,7-Dichloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,

8-Bromo-3-chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 3-Chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 3-Chloro-11-(piperazin-1-yl)-8-trifluoromethyl-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 7-Chloro-2-methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 2-Methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 2-Methyl-11-(piperazin-1-yl)-8-trifluoromethyl-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-4-methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 1,8-Dichloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Bromo-5-methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 7,8-Dichloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 11-(Piperazin-1-yl)-8-trifluoromethyl-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 11-(Piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Fluoro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 11-(Piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine-8-carbonitrile,  
 8-Bromo-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 3-Fluoro-6-piperazin-1-yl-11*H*-dibenzo[*b,e*]azepine,  
 2-(Trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 2-(Trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]oxazepine,  
 8-Chloro-2-(trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5*H*-  
 dibenzo[*b,e*][1,4]diazepine,  
 8-(Trifluoromethanesulfonyloxy)-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 11-(Piperazin-1-yl)-dibenzo[*b,f*][1,4]thiazepin,  
 11-(Piperazin-1-yl)-2,3-dihydro-1,4-benzodioxino[6,7-*b*][1,4]benzothiazepin,  
 8-Chloro-11-[1,4]diazepam-1-yl-5*H*-dibenzo[*b,e*][1,4]diazepine,  
*N*'-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)-*N,N*-dimethyl-ethane-1,2-diamine,  
*N*'-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)-*N,N*-diethyl-ethane-1,2-diamine,  
 8-Chloro-11-(4-methyl-[1,4]diazepam-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-2-methoxy-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
*N*'-(5*H*-Dibenzo[*b,e*][1,4]diazepine-11-yl)-*N,N*-dimethyl-ethane-1,2-diamine,  
 11-[1,4]Diazepam-1-yl-5*H*-dibenzo[*b,e*][1,4]diazepine,  
*N*'-(8-Fluoro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)-*N,N*-dimethyl-ethane-1,2-diamine,  
 8-Fluoro-11-[1,4]diazepam-1-yl-5*H*-dibenzo[*b,e*][1,4]diazepine,





*N*'-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)-*N*-methyl-ethane-1,2-diamine,  
8-Chloro-11-(trans-2,5-dimethyl-piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
8-Chloro-11-(3,5-dimethyl-piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
8-Chloro-11-(3-methyl-piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
8-Chloro-11-(3-phenyl-piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
8-Chloro-5-methyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
8-Chloro-5-benzyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
8-Iodo-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
2-Iodo-8-chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
8-Phenyl-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
8-Chloro-11-(piperidin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
8-Chloro-11-(morpholin-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
5-Allyl-8-chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
6-Chloro-11-(piperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
8-Chloro-5-piperazin-1-yl-11*H*-benzo[*b*]pyrido[2,3-*e*][1,4]diazepine,  
2-Chloro-10-piperazin-1-yl-5*H*-dibenzo[*b,f*]azepin,  
8-Chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]thiazepine,  
8-Chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
8-Chloro-11-(4-methyl-piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
3-Chloro-6-piperazin-1-yl-11*H*-dibenzo[*b,e*]azepine,  
8-Bromo-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
11-(Piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
7-Chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
8-Chloro-3-methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
8-Bromo-3-methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
3-Methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
7-Chloro-3-methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
8-Chloro-4-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
8-Bromo-4-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
4-Methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
2-Bromo-8-chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
2,8-Dibromo-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
2-Bromo-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,

2-Bromo-7-chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 11-(Piperazin-1-yl)-8-trifluoromethyl-dibenzo[*b,f*][1,4]oxazepine,  
 4-Methyl-11-(piperazin-1-yl)-8-trifluoromethyl-dibenzo[*b,f*][1,4]oxazepine,  
 8-Fluoro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Fluoro-3-methoxy-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Fluoro-4-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 2-Bromo-8-fluoro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 3-Methoxy-8-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 4,8-Dimethyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 3-Methoxy-11-(piperazin-1-yl)-8-trifluoromethyl-dibenzo[*b,f*][1,4]oxazepine,  
 2-Bromo-11-(piperazin-1-yl)-8-trifluoromethyl-dibenzo[*b,f*][1,4]oxazepine,  
 6-Chloro-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 2-Bromo-8-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 7-Chloro-4-methyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Phenyl-11-(piperazin-1-yl)-dibenzo[*b,f*][1,4]oxazepine,  
 8-Chloro-11-(piperidin-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine  
 5-Benzyl-8-chloro-11-(piperidin-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Bromo-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one,  
 5,10-Dihydro-dibenzo[*b,e*][1,4]diazepine-11-one,  
 8-Fluoro-5,10-dihydro-dibenzo[*b,e*][1,4]diazepine-11-one,  
 8,5-Dichloro-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-methylsulfanyl-5*H*-dibenzo[*b,e*][1,4]diazepine  
 (8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-(S)-1-pyrrolidin-2-yl-methyl-amine,  
 1-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-piperidine-4-yl-amine,  
 1-(8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-pyrrolidin-3-yl-amine,  
 (8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-(R)-1-pyrrolidin-2-yl-methyl-amine,  
 (8-Chloro-5*H*-dibenzo[*b,e*][1,4]diazepin-11-yl)-pyrrolidin-3-yl-amine,  
 8-Chloro-11-(2,5-diaza-bicyclo[2.2.1]hept-2-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 Acetidin-3-yl-(8-chloro-5*H*-dibenzo[*b,e*][1,4]diazepine-11-yl)amine,  
 7-Bromo-4-(piperazin-1-yl)-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepine,  
 7-Bromo-2-methyl-(piperazin-1-yl)-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepine  
 7-Bromo-2-phenyl-4-(piperazine-1-yl)-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepine,


7-Bromo-10-(piperazin-1-yl)-1,2,3,3a,4,10a-hexahydro-benzo[*b*]cyclopenta[*e*][1,4]diazepine,  
 8-Chloro-11-(4-fluorobenzyl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-(4-fluorophenyl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-(4-nonylphenyl)-5*H*-dibenzo[*b,e*][1,4]diazepine,  
 8-Chloro-11-(pyridin-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine, and  
 8-Chloro-11-(1*H*-pyrazol-4-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine.

59. A method of synthesizing a compound of Formula V or VI,




comprising


reacting a compound of Formula VII



with a compound of Formula VIII




to form a fused ring compound of Formula IX,



and

reacting the compound of Formula IX with a compound of Formula X



to obtain a compound of Formula V,

wherein

X is a halogen;

R<sub>1</sub> is selected from the group consisting of hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, and optionally substituted aryl and arylalkyl;


R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, and R<sub>5</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>,

or R<sub>2</sub> and R<sub>3</sub>, or R<sub>3</sub> and R<sub>4</sub>, or R<sub>4</sub> and R<sub>5</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety;


R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, and R<sub>9</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>,

or R<sub>6</sub> and R<sub>7</sub>, or R<sub>7</sub> and R<sub>8</sub>, or R<sub>8</sub> and R<sub>9</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety.


60. A combinatorial library of at least 220 dibenzo[*b,e*][1,4]diazepine[*a,d*]cycloheptene compounds that can be formed by reacting a compound of Formula VII,



with a compound of Formula VIII and



a compound of Formula X,



wherein

X is a halogen;

R<sub>1</sub> is selected from the group consisting of hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, and optionally substituted aryl and arylalkyl;


R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, and R<sub>5</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>,

or R<sub>2</sub> and R<sub>3</sub>, or R<sub>3</sub> and R<sub>4</sub>, or R<sub>4</sub> and R<sub>5</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety;


R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, and R<sub>9</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>,

or R<sub>6</sub> and R<sub>7</sub>, or R<sub>7</sub> and R<sub>8</sub>, or R<sub>8</sub> and R<sub>9</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety.


61. A combinatorial library of at least 220 dibenzo[*b,e*][1,4]diazepine[*a,d*]cycloheptene compounds that can be formed by reacting a compound of Formula VII,



with a compound of Formula VIII and



a compound of Formula XII,



wherein

X is a halogen;

R<sub>1</sub> is selected from the group consisting of hydrogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>3-8</sub> cycloalkyl, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, and optionally substituted aryl and arylalkyl;

R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, and R<sub>5</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>,

or R<sub>2</sub> and R<sub>3</sub>, or R<sub>3</sub> and R<sub>4</sub>, or R<sub>4</sub> and R<sub>5</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety;

R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, and R<sub>9</sub>, are each independently selected from the group consisting of hydrogen, halogen, optionally substituted C<sub>1-6</sub> alkyl, optionally substituted C<sub>1-6</sub> alkyloxy, optionally substituted C<sub>2-6</sub> alkenyl, optionally substituted C<sub>2-6</sub> alkynyl, optionally substituted C<sub>1-6</sub>-alkoxyalkyl, optionally substituted C<sub>1-6</sub> alkylthio, perhaloalkyl, CN, COR<sub>10</sub>, CONHR<sub>10</sub>, NHCONHR<sub>10</sub>, SO<sub>2</sub>NHR<sub>10</sub>, SO<sub>2</sub>R<sub>10</sub>, OSO<sub>2</sub>R<sub>10</sub>, heteroalkyl, NO<sub>2</sub>, NHCOR<sub>10</sub>,

or R<sub>6</sub> and R<sub>7</sub>, or R<sub>7</sub> and R<sub>8</sub>, or R<sub>8</sub> and R<sub>9</sub> taken together, along with the ring carbons to which they are attached, form a five-membered or six-membered cycloalkyl, heterocyclyl or heteroaryl ring, or a six-membered aryl ring moiety.

62. A pharmaceutical composition comprising a physiologically acceptable carrier, diluent, or excipient, or a combination thereof; and a compound of claim 1.

63. A method of treating a neuropsychiatric disorder comprising administering to said patient a therapeutically effective amount of a compound of claim 1.

64. A method of treating a neuropsychiatric disorder comprising contacting a therapeutically effective amount of a compound of claim 1 with said patient.

65. A pharmaceutical composition comprising a compound of claim 1 and an neuropsychiatric agent.

66. The composition of claim 65, wherein said neuropsychiatric agent is selected from the group consisting of a selective serotonin reuptake inhibitor, norepinephrine reuptake inhibitor, dopamine agonist, muscarinic receptor antagonist, antipsychotic agent, serotonin 2A antagonist, and inverse serotonin 2A agonist.

67. The composition of claim 66, wherein said antipsychotic agent is selected from the group consisting of a phenothiazine, phenylbutylpiperadine, debenzapine, benzisoxidil, and salt of lithium.

68. The composition of claim 67, wherein said phenothiazine is selected from the group consisting of chlorpromazine (Thorazine®), mesoridazine (Serentil®), prochlorperazine (Compazine®), and thioridazine (Mellaril®).

69. The composition of claim 67, wherein said phenylbutylpiperadines is selected from the group consisting of haloperidol (Haldol®), and pimozide (Orap®).

70. The composition of claim 67, wherein said debenzapine is selected from the group consisting of clozapine (Clozaril®), loxapine (Loxitane®), olanzapine (Zyprexa®) and quetiapine (Seroquel®).

71. The composition of claim 67, wherein said benzisoxidil is selected from the group consisting of resperidone (Resperidal®) and ziprasidone (Geodon®).

72. The composition of claim 67, wherein said salt of lithium is lithium carbonate.


73. The composition of claim 66, wherein said antipsychotic agent is selected from the group consisting of Clozaril, Compazine, Etrafon, Geodon, Haldol, Inapsine, Loxitane, Mellaril, Moban, Navane, Orap, Permitil, Prolixin, Phenergan, Reglan, Risperdal, Serentil, Seroquel, Stelazine, Taractan, Thorazine, Triavil, Trilafon, and Zyprexa.

74. The composition of claim 66, wherein said selective serotonin reuptake inhibitor is selected from the group consisting of fluoxetine, fluvoxamine, sertraline, paroxetine, citalopram, escitalopram, sibutramine, duloxetine, and venlafaxine, and pharmaceutically acceptable salts or prodrugs thereof.

75. The composition of claim 66, wherein said norepinephrine reuptake inhibitor is selected from the group consisting of thionisoxetine and reboxetine.

76. The composition of claim 66, wherein said dopamine agonist is selected from the group consisting of sumatriptan, almotriptan, naratriptan, frovatriptan, rizatriptan, zomitriptan, cabergoline, amantadine, lisuride, pergolide, ropinirole, pramipexole, and bromocriptine.


77. The composition of claim 66, wherein said inverse serotonin 2A agonist is the compound of Formula XIII, or a related analog thereof.



(XIII)

78. The composition of claim 66, where in said serotonin 2A antagonist is the compound of Formula XIV, or a related analog thereof:





79. A method of treating neuropsychiatric disorder in a patient comprising administering to said patient a therapeutically effective amount of a composition of claim 65.

80. A method of treating neuropsychiatric disorder in a patient comprising administering to said patient a therapeutically effective amount of a compound of claim 1 and a therapeutically effective amount of a neuropsychiatric agent.

81. The method of claim 80, wherein said administering step comprises administering said compound of claim 1 and said neuropsychiatric agent nearly simultaneously.

82. The method of Claim 80, wherein said administering step comprises administering one of said compound of claim 1 and said neuropsychiatric agent first and then administering the other one of said compound of claim 1 and said neuropsychiatric agent.

83. The method of claim 80, wherein said neuropsychiatric disorder is selected from the group consisting of schizophrenia and related idiopathic psychoses, anxiety, sleep disorders, appetite disorders, affective disorders such as major depression, bipolar disorder, and depression with psychotic features, and Tourette's Syndrome, drug-induced psychoses, psychoses secondary to neurodegenerative disorders such Alzheimer's or Huntington's Disease.